

Do not re-use container. Rinse equipment and containers and bury with waste in a location away from water supplies. Never use container in any connection with food, feed or drinking water.

This product is toxic to fish, birds, and other wildlife. Apply only as specified on this label. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not use on any plants to be used for food or feed.

#### DIRECTIONS FOR USE

Zectran controls a wide range of pests, including snails, slugs and many insects on turf, shade trees, flowers and shrubs.

Apply Zectran as a spray, diluted with water, using 3 teaspoonfuls per gallon (1½ qts. per 100 gal. water). Applications may be made with hand sprayers, power equipment, and on turf with hose attached sprayers. Use enough spray to give thorough coverage.

**ORNAMENTALS:** Can be used on many plants including most annual and perennial flowers, bedding plants, ground covers, shrubs, evergreens and trees,

**S A BRAND 5 O** ®

# ZECTRAN\*2E

**INSECTICIDE FOR ORNAMENTALS AND TURF.**

Use 3 teaspoonfuls per gallon.

**ACTIVE INGREDIENTS:**

4-(dimethylamino)-3,5-xylol methylcarbamate	22.0%
Aromatic petroleum derivative solvent	31.0%
<b>INERT INGREDIENTS</b>	<b>47.0%</b>
	<b>100.0%</b>

Contains 2 lbs. 4-(dimethylamino)3,5-xylol methylcarbamate per gallon.

\*ZECTRAN, trademark of The Dow Chemical Co.  
U.S. Patent No. 3,084,098

#### KEEP OUT OF REACH OF CHILDREN

**WARNING:** May be harmful or fatal if swallowed. May cause irritation. Avoid eye and skin contact. Do not breathe spray mist. Do not contaminate food, feed stuffs and domestic water supplies. Wash thoroughly after handling. Flush contaminated eyes with plenty of water and get medical attention. If swallowed call a physician immediately.

**NOTE TO PHYSICIAN:** Active ingredient has cholinergic effects. Treat symptomatically. Atropine is an antidote.

Packaged and Distributed by

**Southern Agricultural Insecticides, Inc.**

Palmetto, Fla. Hendersonville, N. C. Boone, N. C.

such as ash, azalea, camellia, elm, honeysuckle, juniper, lilac, locust, maple, pines, privet, rose and spruce.

For the control of armyworms, leaf hoppers, leaf miners, mealybugs, mites, scale crawlers, spruce budworms, tent caterpillars and thrips.

**TURF:** For use on Bermuda grass, blue grass, dichondra and most other turf grasses and ground covers. For control of cutworms, millipeds, pillbugs, sod webworms, snails and slugs. Water turf thoroughly before treating. Spray infested area using 1 gallon of diluted spray per 130 to 180 square feet of area. Soak the debris and soil under infested plants. Repeat as necessary.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**NOTE:** Do not use on Maidenhair fern, cape chestnut, or Geraldton wax flower. Excessive dosage may cause leaf injury to new growth on roses, phlox and petunias.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.

EPA Reg. N. 829-207

Net Contents Liquid:

**ACCEPTED**

SEP 13 1972

FEDERAL INSECTICIDE  
AND RODENTICIDE ACT  
POISON REGISTERED  
EPA REG. NO. 829-207