


ACCEPTED
 MARCH 2, 1972
 921-211



SA-50 BRAND Net Wt. _____ Lbs.

BALFIN GRANULES

FOR LAWNS AND GOLF COURSES

For Use on Fescue, Blue, Bermuda, Centipede, Bahia,
St. Augustine and Zoysia Grasses

For the pre-emergence control of smooth and hairy Crabgrass, Water Grass (Barnyard Grass), Yellow and Green Foxtail, Goosegrass, and certain other weeds from seed

ACTIVE INGREDIENTS:	
N-butyl-N-ethyl-3,4-difluoro-2,6-dinitro-P-toluidine	2.0%
INERT INGREDIENTS	98.0%
Balafin - the Trademark of Elanco Products Co. Benefin	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION!

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact flush with water. Balfin Granules can stain skin and clothing. Wash thoroughly with soap and water after using. Do not contaminate feed or foodstuffs. Destroy empty container and bury with waste Balfin Granules in an isolated area. Do not graze or feed forage from treated areas to livestock. Toxic to fish. Do not contaminate any body of water by application, cleaning of equipment or disposal of waste. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.
 PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.

Recommendations for Use in the Southern Appalachian and Piedmont Plateau Areas

Apply SA-50 Balfin Granules to established turfs before germination of annual grass weeds. (Preferably before March 25th in Piedmont and April 10th in Mountain areas)

Use 1½ lbs. per 1000 sq. ft. for fine-leaved fescues and not less than 2 lbs. per 1000 sq. ft. and not more than 3½ lbs. per 1000 sq. ft. for all other recommended turfs. Apply with a spreader designed to apply granular materials that has been calibrated before using. Apply to turf, at indicated rates evenly without skips or overlaps. Water thoroughly after application. Application may be repeated in 5 months to inhibit late summer germination of grass weeds.

PRECAUTIONS: Do not apply in spring to Blue Grass planted the previous fall. Do not apply to Bent or Dichondra turfs. Do not apply to newly sprigged St. Augustine, Bermuda, Centipede or Zoysia turfs until these turfs are well established as root inhibition may result. Applications in excess of 1½ lbs. per 1000 sq. ft. may thin fine-leaved fescues. Do not reseed or overseed within 6 weeks before or after application of Balfin Granules. Do not apply to turf that is under stress from drought, nematode infestation or stress from other causes. Granules may stain masonry. Remove granules from walkways, flagstones, etc. immediately following application.

Spreader should be calibrated before using

TO CALIBRATE: Measure the spread width (swath of the spreader). Place 1 lb. in spreader and apply in a straight line. Determine the distance traveled to deliver the pound of granules. Adjust the spreader setting and repeat the process until the spreader will deliver 1 lb. over the distance indicated in the chart for the measured swath width and the desired dosage in lbs. per 1000 sq. ft. of turf.

Swath Width	Feet of Travel at Indicated Dosage of Spreader		
	1½ Lbs.	2 Lbs.	3½ Lbs.
1 foot	170 ft.	84 ft.	48 ft.
5 feet	33 ft.	100 ft.	57 ft.
4 feet	37 ft.	125 ft.	72 ft.
3 feet	47 ft.	150 ft.	91 ft.

A
M
UNDE
FUNG
FOR E
ED U
T

SA-50[®] BRAND

Net Wt. _____ Lbs.

BALFIN[®] GRANULES

FOR LAWNS AND GOLF COURSES

**For Use on Fescue, Blue, Bermuda, Centipede, Bahia,
St. Augustine and Zoysia Grasses**

For the pre-emergence control of smooth and hairy Crabgrass, Water Grass (Barnyard Grass), Yellow and Green Foxtail, Goosegrass, and certain other weeds from seed.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl,a,a,a-trifluoro-2, 6-dinitro-P-toluidine 2.0%

INERT INGREDIENTS 98.0%

Balafin[®] the Trademark of Elanco Products Co. Benefin 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION! Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of contact flush with water. Balfin Granules can stain skin and clothing. Wash thoroughly with soap and water after using. Do not contaminate feed or foodstuffs. Destroy empty container and bury with waste Balfin Granules in an isolated area.

Do not graze or feed forage from treated areas to livestock.

Toxic to fish. Do not contaminate any body of water by application, cleaning of equipment or disposal of waste.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. Buyer assumes all risks of use, storage or **handling of this material not in strict accordance with the directions and precautions given herewith.**

Southern Agricultural Insecticides, Inc.

PALMETTO, FLA. BOONE, N. C. HENDERSONVILLE, N. C.