

SA BRAND 50[®]

7½% Captan Fungicide Dust

ACTIVE INGREDIENTS:

Captan (N-[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide) 7.5%

INERT INGREDIENTS 92.5%

U. S. Pat. Nos. 2,553,770; 2,553,771; 2,553,776 **100.00%**

— SUGGESTED FOR USE ON —

STRAWBERRIES: Botrytis Rot, 30 to 40 lbs. per acre. Make applications when new growth starts in spring before fruit starts to form. Repeat applications weekly. Severe Infection Conditions, continue through harvest period, treating immediately after each picking.

SOIL AND GREENHOUSE BENCH TREATMENT: Use 2-2½ lbs. per 1000 sq. ft. (100 lbs. per acre). Work into upper 3 or 4 inches of soil before planting. For damp-off and root rot diseases of seedlings and transplants of roses, other shrubs, flowers, lawn seedbeds, and vegetables such as beans, peas, celery, cabbage, eggplant, pepper, and tomato.

ORNAMENTALS: Rose, black spot; Carnation, rust, leaf spot; Chrysanthemums, flower blight — Apply at first sign of disease to given even coverage of upper and lower leaf surfaces. Repeat at 7 to 10-day intervals as necessary. Shorten interval during periods of frequent rain and heavy dew.

NOTE: This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Keep Out of Reach of Children

CAUTION! Do not breathe dust. Avoid contact with skin and eyes. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Store in cool, dry place. Protect from excessive heat.

Do not reuse container. Destroy when empty.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with the directions and precautions given herewith.

Southern Agricultural Insecticides, Inc.

Palmetto, Florida Boone, N. C. Hendersonville, N. C.

EPA Reg. No. S29-215

A

NET WT.

