

7 1/2% CAPTAN FUNGICIDE DUST

per application
829-215

ACTIVE INGREDIENTS:

*Captan
Related Derivatives

INERT INGREDIENTS

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

BY WT.

7.8%
0.2%
92.5%

**KEEP OUT OF REACH OF CHILDREN
DANGER — AVISO**
CAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**
DANGER
Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.
STATEMENT OF PRACTICAL TREATMENT
IF IN EYES: Flush with plenty of water. Call a physician.
IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.
IF INHALED: Remove victim to fresh air if not breathing give artificial respiration preferably mouth-to-mouth. Get medical attention.
ENVIRONMENTAL HAZARDS
This product is toxic to fish. Do not apply directly to water or wetlands, swamps, bays, streams, and potholes. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

NET WEIGHT: 50 POUNDS

EPA EST NO 829-FL-1 (3), 329-NC-1 (4), 829-NC-2 (5) Number following Est. No. corresponds to first digit in the Lot No. on bottom of bag

Southern Agricultural Insecticides, Inc.
Palmatto, FL 34220 Hendersonville, NC 28753 Boone, NC 28607
EPA Reg. No. 829-215 AA

DIRECTIONS FOR US
It is a violation of Federal law to use this product in a manner inconsistent with its labeling
READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

REENTRY STATEMENT
Keep all unprotected persons out of operating areas or vicinity where there may be drift. Do not allow persons to enter treated areas within 4 days following application unless a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Conspicuously post reentry information at site of application.

Certain states may require more restrictive reentry intervals, consult your State Department of Agriculture for further information. Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

DIRECTIONS
Mixers, loaders, when mixing and loading this product, must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves if entering treated area within 4 days following application. Leather or fabric gloves are not acceptable. A dust mask must be worn when mixing and loading this product. During application only, if from a tractor with a completely enclosed cab, a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab and worn while exiting.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

NOTE: Do not apply this product by air.

ORNAMENTALS: ROSES - Black Spot. CARNATIONS - Rust, Leaf Spot. CHRYSANTHEMUMS - Flower Blight — Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for Damp Off and Root Rot Disease on Seedling or Transplants of Roses (and other shrubs and trees), Many Flowers and Lawn Seedbeds — 1.75 to 2.3 lbs per 1000 sq ft (75 to 100 lbs per acre). Work into upper 3 to 4 inches of soil before planting. Only the applicator is permitted to be in the greenhouse during application of captan. Open vents to greenhouse during application and for at least 1 hour after application.

Workers planting in captan treated soil in greenhouses must wear chemical resistant gloves.

STRAWBERRIES: Botrytis Rot — Use 40 lbs per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat applications weekly. Severe Infection Conditions — Continue through harvest period, treating immediately after each picking.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Store in cool, dry place. Protect from excessive heat. For help with any spill, leak, fire or exposure involving this material, call day or night (415) 233-7737.

PESTICIDE DISPOSAL

This product is acutely hazardous. Improper disposal of excess pesticide, mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities by burning. If burned, stay out of smoke.

CONDITIONS OF SALE: 1. Southern Agricultural Insecticides, Inc warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Southern Agricultural Insecticides, Inc. neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Southern Agricultural Insecticides, Inc. is not liable for any actual and unforeseeable factors beyond Southern Agricultural Insecticides, Inc.'s control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Southern Agricultural Insecticides, Inc. under 1 above) resulting from the handling, storage, and use of this material.

ORM-E

**RQ HAZARDOUS
SUBSTANCE, SOLID,
N.O.S., NA 9188,
(Captan)**

