

ACCEPTED

Handwritten signature

829-217

DIRECTIONS

Benomyl Fungicide is recommended for the control of certain diseases of roses, flowers, ornamentals, and turf. It is highly effective for the control of many fungus diseases, providing both curative (eradicant) action and protective action. Benomyl Fungicide is a wettable powder to be mixed with water for application as a spray. Addition of a spreader-sticker such as SA-50 Brand Polyethylene Spreader-sticker to the spray mixture enhances curative action of the fungicide and improves distribution of the spray on hard-to-wet plants such as roses.

ROSES — Powdery Mildew, Black Spot: Use one tablespoonful per 2 gals. of water. Begin application when disease first appears and repeat at 10 to 14 day intervals throughout the growing season; shorten intervals during humid, rainy weather.

FLOWERS, ORNAMENTALS such as aster, chrysanthemum, snapdragon, crepe myrtle, azalea, gardenia and hydrangea — **Powdery Mildew, Botrytis, Grey Mold:** Use one tablespoonful per 2 gals. of water. Begin application when disease first appears and repeat at 10 to 14 day intervals as long as needed.

BULB DIP — Tulip, Iris, Easter Lily, Daffodil, and Gladiolus Bulbs and corms for control of Fusarium basal rot and Penicillium infections. Dip clean bulbs in a warm water (80-85°F) suspension of Benomyl Fungicide (2 tablespoonfuls per gallon) for 30 minutes, within 48 hours after digging. Maintain suspension with sufficient agitation. Dry bulbs after treatment for storage.

Benomyl Fungicide

BRAND

ACTIVE INGREDIENT

Benomyl {Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate}	50%
INERT INGREDIENTS	50%
	<hr/> 100%

U.S. Pat Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION: MAY IRRITATE EYES, NOSE, THROAT, AND SKIN

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. Keep away from fire or sparks.

In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

IMPORTANT—Never allow Benomyl Fungicide to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness. Keep container closed when not in use. Do not re-use container; destroy when empty.

TURF — Dollar Spot (Sclerotinia) Large Brown Patch (Rhizoctonia) at the rate of 2 oz. (12 tablespoons) per 1000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gals. per 1000 sq. ft. at the earliest appearance of disease and repeat at 10 to 14 day intervals as long as needed. Under unusually favorable conditions are unusually favorable for control of disease, reduce interval to 5 to 7 days. Do not graze or feed clippings from treated areas to livestock.

This product is toxic to fish. Keep away from streams or ponds. Do not apply where drift is likely to occur. Do not apply when conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product as specified on this label.

Buyer assumes all risk of use. Handling not in strict accordance with directions and precautions given hereon.

Packaged and Distributed by

Southern Agricultural Insect

Palmetto, Fla. Hendersonville, N.C.

EPA Reg. No. 829-217 Net

ACCEPTED

December 1972

829 217



BRAND

Benomyl Fungicide

ACTIVE INGREDIENT

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate] 50%

INERT INGREDIENTS 50%
100%

U.S. Pat Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION: MAY IRRITATE EYES, NOSE, THROAT, AND SKIN

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. Keep away from fire or sparks.

In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention.

IMPORTANT—Never allow Benomyl Fungicide to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness. Keep container closed when not in use. Do not re-use container; destroy when empty.

TURF — Dollar Spot (*Sclerotinia homeocarpa*), **Large Brown Patch** (*Rhizoctonia solani*): Apply at the rate of 2 oz. (12 tablespoons) per 1000 sq. ft. in sufficient water to obtain thorough coverage, usually 2 to 5 gals. per 1000 sq. ft. Apply at earliest appearance of disease and continue at 10 to 14 day intervals as long as needed. When conditions are unusually favorable for development of disease, reduce interval to 5 to 7 days. Do not graze or feed clippings from treated areas to livestock.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Buyer assumes all risk of use, storage, or handling not in strict accordance with the directions and precautions given herewith.

Packaged and Distributed by

Southern Agricultural Insecticides, Inc.

Palmetto, Fla. Hendersonville, N.C. Boone, N.C.

EPA Reg. No. 829-217

Net Weight 6 oz.

ended for the
flowers, orna-
effective for the
providing both
protective action.
powder to be
on as a spray.
such as SA-50
er to the spray
of the fungi-
of the spray on

Spot: Use one
iter. Begin ap-
ears and repeat
out the growing
humid, rainy

as aster, chrys-
myrtle, azalea,
powdery Mildew,
tablespoonful per
on when disease
14 day intervals

Lily, Daffodil,
for control of
m infections.
(80-85°F) sus-
2 tablespoonfuls
48 hours after
sufficient agi-
for storage.