

DIRECTIONS FOR USE

Chlorothalonil Fungicide is highly effective for the control of a number of important plant diseases. It is compatible with wettable powder formulations of most commonly used fungicides and insecticides. Do not combine with any material unless prior use has shown the combination compatible.

LAWNS AND ORNAMENTAL TURF

Helminthosporium leaf spot and melting out, Curculiaris leaf spot, Dollar spot, Brown patch, Gray leaf spot, Stem rust of Blue Grass and Copper spot. Apply 2 to 6 ounces in sufficient water (5 to 15 gal.) to cover 1,000 sq. ft. Do not mow or water for at least 24 hours after treatment. Treat grass at 7 to 14 day intervals as long as needed. Under severe disease conditions use higher rate and treat every 7 days. Do not graze treated areas or feed clippings to livestock.

ORNAMENTALS

Begin applications at the first signs of disease. Use 1 1/2 Tablespoonsful per gallon of water. Repeat at 7 to 14 day intervals as long as needed on the following plants for control of listed diseases.

Rose—Black spot, Blossom blight.

Chrysanthemum—Ascochyta ray blight, Gray mold.

Geranium—Botrytis blight.

Zinnia—Powdery mildew.

VEGETABLES

Begin applications at the first sign of disease or when disease first threatens. Use 2 to 3 Tablespoonsful per gallon of water. Repeat at 7 to 10

S A 5 O[®] BRAND LAWN ORNAMENTAL & VEGETABLE FUNGICIDE

ACTIVE INGREDIENT:
Chlorothalonil (tetrachloroisophthalonitrile)* 75%
INERT INGREDIENTS 25%
100%

Diamond Shamrock Chemical Co. U.S. Pat. No. 3,290,353 & 3,331,735.

CAUTION: Keep Out Of Reach Of Children

Do not take internally. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Can be irritating to eyes or skin. Wash thoroughly with soap and warm water after handling. In case of contact with eyes, flush with plenty of water for at least 15 minutes.

Packaged and Distributed by

Southern Agricultural Insecticides, Inc.

Palmetto, Florida Hendersonville, N.C. Boone, N.C.
EPA Reg. No. 829-232 Net Weight 6 oz.

day intervals or as necessary to maintain control on the following crops for control of listed diseases. May be used up to the day of harvest except on Snap Beans.

Snap Beans—Rust. Do not use within 7 days of harvest. Do not graze treated area or feed treated plant parts to livestock.

Cabbage, Cauliflower, Broccoli and Brussels Sprouts—Alternaria leaf spot (Southeast only), Downy mildew.

Carrot—Early and Late Blight.

Cucumber, Cantaloupe, Muskmelon, Honeydew, Watermelon, Squash and Pumpkin—Anthracnose, Downy mildew, Gummy stem blight, Leaf blight and Scab.

Potato—Early and Late Blights, Botrytis Vine Rot.
Tomato—Early and Late Blights, Gray Leaf Spot, Gray Leaf Mold, Rhizoctonia fruit rot, Botrytis Gray Mold, Anthracnose.

NOTE: Chlorothalonil may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with anti-histamines or steroid creams and/or systemic steroids.

Do not store near feed or foodstuffs. Do not reuse container. Destroy when empty.

This product is toxic to fish. Keep out of lakes, ponds, and streams. Do not contaminate water by cleaning equipment or disposal of wastes. Apply only to areas specified on label.

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with the directions and precautions given herewith.

