

Final 152

655 / 164350
17 / 1

Southern Agricultural Insecticides, Inc.
P.O. Box 218
Palmetto, FL 33561-0218

Attention: Ronald L. Costa Rica

Gentlemen:

JAN 24 1986

Subject: Lawn Ornamental Vegetable Fungicide
EPA Registration No. 829-232
Your Letter Dated November 25, 1985

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted at the next printing.

1. Change "Caution" to "Danger." Justification or citation of specific toxicology data must be provided in order to consider an alternative hazard signal warning word other than "Danger."
2. Add "Skin Sensitizer" to the Precautionary Statements.

A stamped copy of one label is enclosed for your records.

Sincerely yours,

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosure

88852:Noles;J-1;KENCO:1/13/86:1/22/86:ab:vo
CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

5A-50

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 Corrosive, causes severe eye damage. Do not take internally. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist.

ENVIRONMENTAL HAZARDS
 This product is toxic to fish, aquatic invertebrates, and marine estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE
 IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
 Chlorothalonil Fungicide is highly effective for the control of a number of important plant diseases. It is compatible with wettable powder formulations of most commonly used fungicides and insecticides. Do not combine with any material unless prior use has shown the combination compatible.

NOTE TO USER: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

LAWNS AND ORNAMENTAL TURF
 Helminthosporium leaf spot and melting out, Curvularia leaf spot, Dollar spot, Brown patch, Gray leaf spot, Stem rust of Blue Grass and Copper spot. Apply 2 to 6 ounces in sufficient water (5 to 15 gal) to cover 1,000 sq ft. Do not mow or water for at least 24 hours after treatment. Treat grass at 7 to 14 day intervals as long as needed. Under severe disease conditions use higher rate and treat every 7 days.

Do not graze treated areas or feed clippings to livestock.

ORNAMENTALS
 Begin applications at the first sign of disease. Use 1½ Tablespoonsful per gallon of water. Repeat at 7 to 14 day intervals as long as needed on the following plants for control of listed diseases:
 Rose—Black spot, Blossom blight
 Chrysanthemum—Ascochyta ray blight, Gray mold
 Geranium—Botrytis blight
 Zinnia—Powdery mildew

VEGETABLES
 Begin applications at the first sign of disease or when disease first threatens. Use 2 to 3 Tablespoonsful per gallon of water. Repeat at 7 to

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 Corrosive, causes severe eye damage. Do not take internally. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist.

ENVIRONMENTAL HAZARDS
 This product is toxic to fish, aquatic invertebrates, and marine estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE
 IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.
 Chlorothalonil Fungicide is highly effective for the control of a number of important plant diseases. It is compatible with wettable powder formulations of most commonly used fungicides and insecticides. Do not combine with any material unless prior use has shown the combination compatible.

NOTE TO USER: Wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

LAWNS AND ORNAMENTAL TURF
 Helminthosporium leaf spot and melting out, Curvularia leaf spot, Dollar spot, Brown patch, Gray leaf spot, Stem rust of Blue Grass and Copper spot. Apply 2 to 6 ounces in sufficient water (5 to 15 gal) to cover 1,000 sq ft. Do not mow or water for at least 24 hours after treatment. Treat grass at 7 to 14 day intervals as long as needed. Under severe disease conditions use higher rate and treat every 7 days.

Do not graze treated areas or feed clippings to livestock.

ORNAMENTALS
 Begin applications at the first sign of disease. Use 1½ Tablespoonsful per gallon of water. Repeat at 7 to 14 day intervals as long as needed on the following plants for control of listed diseases:
 Rose—Black spot, Blossom blight
 Chrysanthemum—Ascochyta ray blight, Gray mold
 Geranium—Botrytis blight

**LAWN
 ORNAMENTAL
 VEGETABLE FUNGICIDE**

ACTIVE INGREDIENT:
 Chlorothalonil (tetrachloroisophthalonitrile) 75%
INERT INGREDIENTS 25%
TOTAL 100%
 *SDS BIOTECH CORP
 U.S. PATENT NO 3,290,353 and 3,331,735

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**
STATEMENT OF PRACTICAL TREATMENT
 IF IN EYES: Flush eyes with water for 15 minutes, get medical attention immediately. IF ON SKIN: Wash exposed areas of skin with soap and warm water. IF INHALED: Remove person to fresh air and get medical attention immediately.
NOTE TO PHYSICIAN: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.
 See side panel for additional precautionary statements.

EPA EST NO 829-NC-1 (3), 829-NC-1 (4), 829-NC-2 (6) Number following Est. No. corresponds to first digit in Lot No. on side of label.
Southern Agricultural Insecticides, Inc.
 Palmetto, FL Hendersonville, NC Boone, NC
 EPA Reg No 829-232 AA Net Weight: 1 Lb

**ACCEPTED
 with COMMENTS
 in EPA Letter Dated**

**LAWN JAN 24 1986
 ORNAMENTAL
 VEGETABLE FUNGICIDE**

ACTIVE INGREDIENT:
 Chlorothalonil (tetrachloroisophthalonitrile) 75%
INERT INGREDIENTS 25%
TOTAL 100%
 *SDS BIOTECH CORP
 U.S. PATENT NO 3,290,353 and 3,331,735

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**
STATEMENT OF PRACTICAL TREATMENT
 IF IN EYES: Flush eyes with water for 15 minutes, get medical attention immediately. IF ON SKIN: Wash exposed areas of skin with soap and warm water. IF INHALED: Remove person to fresh air and get medical attention immediately.
NOTE TO PHYSICIAN: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.
 See side panel for additional precautionary statements.

EPA EST NO 829-FL-1 (3), 829-NC-1 (4), 829-NC-2 (6) Number following Est. No. corresponds to first digit in Lot No. on side of label.
Southern Agricultural Insecticides, Inc.

10 day intervals or as needed for control of listed diseases except on Snap Beans.
 Snap Beans—Rust Do not treat area or feed treated Cabbage, Cauliflower, Broccoli spot (Southeast only) CARROT—Early and Late Cucumber, Cantaloupe, Squash and Pumpkin—Anthracnose, Leaf blight and Scab Potato—Early and Late Blight Tomato—Early and Late Blight
 Rhizoctonia fruit rot, Botrytis
NOTE TO USER: This product effects characterized by redness and redness or rash on exposed skin should contact a physician.

STORAGE
 Pesticide wastes are toxic. Pesticide spray or rinse water cannot be disposed of. Contact your State Agency, or the Hazardous Waste Regional Office for guidance. Store pesticide in the original container. To dispose of pesticide, see layers of newspaper and dispose thoroughly before disposal.

WARRANTY AND LIMITATION
 SELLER warrants that the description and is reasonably intended in accordance with directions. BUYER assumes the risk of use. MAKES no other express or implied warranty. No agent of SELLER is authorized to make any specific reference to this warranty or to which a claim is made.

10 day intervals or as needed for control of listed diseases except on Snap Beans.
 Snap Beans—Rust Do not treat area or feed treated Cabbage, Cauliflower, Broccoli spot (Southeast only) CARROT—Early and Late Cucumber, Cantaloupe, Squash and Pumpkin—Anthracnose, Leaf blight and Scab Potato—Early and Late Blight Tomato—Early and Late Blight
 Rhizoctonia fruit rot, Botrytis
NOTE TO USER: This product effects characterized by redness and redness or rash on exposed skin should contact a physician.

STORAGE
 Pesticide wastes are toxic. Pesticide spray or rinse water cannot be disposed of. Contact your State Agency, or the Hazardous Waste Regional Office for guidance. Store pesticide in the original container. To dispose of pesticide, see layers of newspaper and dispose thoroughly before disposal.

WARRANTY AND LIMITATION
 SELLER warrants that the description and is reasonably intended in accordance with directions. BUYER assumes the risk of use. MAKES no other express or implied warranty.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 829-232