

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician.
IF SWALLOWED: Call a physician immediately. Induce vomiting immediately by giving the victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear long sleeve shirt, long pants, shoes & socks, and goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to terrestrial areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply the product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with **DIRECTIONS FOR USE** and **PRECAUTIONARY STATEMENTS** stated on this label.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

5A-60 Broad Fruit Spray Concentrate is formulated for use as a general purpose spray to control the insects and diseases listed for the fruit crops given below. Certain insects or diseases may not be controlled or may be difficult to control and an alternative pesticide for specific pests should be used. Application of a fungicide alone, closer to harvest than this product may be used, is often advisable. Check on current recommendations for a suitable material.

Mix this material at the rate of 8 tablespoons per gallon of water (10 ounces makes 6 gallons of spray). Good agitation is essential since the material will settle out of spray water. Apply solution at the rates given for each type of fruit.

Interval in days, between last application of this product and harvest is shown in () parenthesis after each crop.

NOTE: Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse.



Home & Garden

FRUIT SPRAY CONCENTRATE

ACTIVE INGREDIENTS:	By Wt.
Methidathion, 0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate	6.0%
Methoxychlor technical	9.0%
Equivalent to 7.92% 2,2-bis(4-methoxyphenyl)-1,1,1-trichloroethane and 1.08% other isomers and reaction products	
Captan	5.84%
Related Derivatives	0.16%
Surf	25.0%
INERT INGREDIENTS:	54.0%
TOTAL:	100.0%
N-t-butylbenzothiazol-4-ylcyclohexane-1,2-dicarboximide	

Keep Out of Reach of Children
DANGER

See Side Panel for Additional Precautionary Information



Southern Agricultural Insecticides, Inc.

Palm Beach, FL 33422 Hendersonville, NC 28752 Boone, NC 28607
 J4 EPA Reg. No. 829-236

Net Weight: 5 lbs. (2.27 kg.)

Wear chemically resistant gloves when using. When using wear long pants and long sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before reusing.

APPLES (17): Aphids, Mites, Bud Moth, Codling Moth, Plum Curculio, Red-Banded Leaf Roller, Apple Maggot, Japanese Beetle, Tent Caterpillars, Scab, Black Rot, Blister Rot, White Rot, Sooty Blotch, Froggyleaf Spot, and Powdery Mildew. Apply solution at the rate of 1 - 2 pints per tree on small trees, 1 - 2 quarts on medium trees, and 1 - 3 gallons on large mature trees. Begin applications when new spring growths appear and repeat at 6 to 7 day intervals up to bloom. After petal fall repeat at 7 to 14 day intervals.

CHERRIES (7): Aphids, Bud Moth, Cherry Fruit Flies, Fruit Tree Roller, Cherry Fruitear, Japanese Beetle, Plum Curculio, Rose Chafer, Tent Caterpillars, Carliarworms, Botrytis Rot, Brown Rot, Leaf Spot, Powdery Mildew, Rust and Scab. Apply 1 - 2 pints of solution to small trees, 1 - 2 quarts to medium trees, and 4 - 5 quarts to large mature trees. Begin applications during bloom. Repeat at petal fall and at 10 - 14 day intervals thereafter.

GRAPES (14): Aphids, Mites, Mealybugs, Leafhoppers, Berry Moth, Japanese Beetle, Leaf Skeletonizer, Rose Chafer, Downy Mildew, Black Rot and Powdery Mildew. Apply 3 quarts of solution to 100 sq. ft. of vine. Begin applications at first signs of new spring growth and repeat at 10 - 14 day intervals.

PEACHES (21): Aphids, Mites, Oriental Fruit Moth, Plum Curculio, Japanese Beetle, Rose Chafer, Tent Caterpillar, Carliarworm, Brown Rot, Fungus, Scab, Leafspot, Rust and Powdery Mildew. Apply 1 - 2 pints of solution to small trees, 1 - 2 quarts on medium trees, and 4 - 5 quarts on large mature trees. Begin applications just prior to blossom opening, repeat during bloom, when petals have fallen, and at 7 - 14 day intervals thereafter.

STRAWBERRIES (3): Aphids, Spider Mite, Strawberry Weevil, Flea Beetles, Spittlebugs, Botrytis Rot, Leaf Spots and Powdery Mildew. Apply 1 gallon of solution to 100 sq. ft. of bed area. Begin application when new growth starts and repeat at 7 to 10 day intervals.

STORAGE AND DISPOSAL

Store only in original container in a cool dry place inaccessible to children and pets and away from food and feedstuffs. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse container. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with **DIRECTIONS FOR USE** and **PRECAUTIONARY STATEMENTS** stated on this label.

EPA EST No. 829-FL-1 (3), 829-NC-1 (4), 829-NC-2(5) Number following establishment No. corresponds to first digit in lot number.

ACCEPTED
 S 28-97
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this insecticide is registered.
 829-236

PM-04
 05/10/1997 12:44
 941-723-2974
 829-236
 BEST COPY AVAILABLE
 SOUTHERN AG FLA
 8/28/97
 PAGE 02
 191