

S A  
BRAND

# DIAZINON 14G

5 0°

## Granular Insecticide

ACCEPTED

ACTIVE INGREDIENT:  
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate

INERT INGREDIENTS:

Diazinon® trademark of CIBA-GEIGY

KEEP OUT OF REACH

**CAUTION:** Harmful if swallowed. Avoid contamination of food and feed. Do not eat, drink, or use food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. May be absorbed through skin.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy it by burying or burning. Avoid smoke or fumes.

Follow directions carefully. Timing, method of application, weather conditions and other factors influencing the effectiveness of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.

### DIRECTION FOR USE

Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, harrow, or other suitable means for greatest effectiveness.

#### Tobacco

Cutworms: Broadcast 14-21 lbs. per acre before transplanting. Work into soil 2-4 inches.

Mole Crickets: Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting, and work into soil 1-2 inches.

Wireworms. Broadcast 14-21 lbs. per acre 3-4 weeks before transplanting, and work into soil 6-9 inches.

#### VEGETABLE CROP INSECTS

Broccoli, Brussels Sprouts, Cabbage, and Cauliflower

Root Maggots: Broadcast 14-21 lbs. per acre before planting or transplanting. Work into soil 3-4 inches.

#### Onions

Onion Maggots: Apply 14-28 lbs. per acre as a broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable means.

EPA Reg. No. 829-237

Net Weight: 5 Lbs.

PACKAGED AND DISTRIBUTED BY

**Southern Agricultural Insecticides, Inc.**

PALMETTO, FLA. HENDERSONVILLE, N.C. BOONE, N.C.

EPA EST 829-F1-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No. on bottom of bag.

#### OTHER VEGETABLE INSECTS

##### Cutworms (Surface and Subterranean)

Broccoli	Endive (escarole)	Pole Beans
Brussels Sprouts	Kale	Snap Beans
Cabbage	Lettuce	Spinach
Cantaloupes	Lima Beans	Summer Squash
Cauliflower	Muskmelons	Swiss Chard
Celery	Farsley	Tomatoes
Collards	Peas	Watermelons
Cucumbers	Peppers	Winter Squash

Broadcast 14-28 lbs. per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms, 3-6 inches to control subterranean cutworms.

Potatoes: Broadcast 14-28 lbs. per acre before planting. Work into soil 4-6 inches.

##### Mole Crickets

Broccoli	Endive (escarole)	Peppers
Brussels Sprouts	Kale	Radishes
Cabbage	Lettuce	Tomatoes
Cauliflower	Mustard	Turnips
Collards		

Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

##### Wireworms

Broccoli	Endive (escarole)	Pole Beans
Brussels Sprouts	Kale	Snap Beans
Cabbage	Lettuce	Spinach
Cantaloupes	Lima Beans	Summer Squash
Carrots	Muskmelons	Sweet Potatoes
Cauliflower	Onions	Swiss Chard
Celery	Farsley	Tomatoes
Collards	Peas	Watermelons
Cucumbers	Peppers	Winter Squash

Broadcast 21-28 lbs. per acre just prior to planting. Work into soil 4-8 inches.

Potatoes: Broadcast 21-42 lbs. per acre before planting. Work into soil 4-8 inches or apply 7-21 lbs. per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

Sweet Potatoes (Southeastern U.S.): Broadcast 21 lbs. per acre over tops of vines when roots begin to enlarge.

#### Equivalents for small area treatment.

Lbs. per acre	7	14	21	28	42
Lbs. Per 1,000 sq. ft.	1/6	1/3	1/2	2/3	1

#### LAWN INSECTS

For control of Chinch Bugs, Earwigs, Ticks, Ants, Lawn Moths, (Sod Webworms), Bermuda Mites, Crickets, Armyworms, Brown Dog Ticks, Chiggers, Clover Mites, Cutworms, Fleas, Le-shoppers, Sowbugs, Millipedes, Springtails (Collembola) apply 1/4 - 1 lb. per 1,000 sq. ft. For longer residual control and for use on dense turf such as St. Augustine Grass, use 1-1 1/4 lbs. per 1,000 sq. ft.

Apply granules with fertilizer spreader. Apply when grass is dry. Brush lawn to work granule down to soil surface and water lawn thoroughly after application. Repeat application when necessary.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

JUN 14 1977  
JUN 14 1977  
100.0%  
86.0%

829-237

6-14-77

1021