



CUTWORM & CRICKET BAIT

FOR VEGETABLES, FLOWERS, FRUITS,
LAWNS & ORNAMENTAL PLANTS

FOR THE CONTROL OF CUTWORMS, ARMYWORMS,
FIELD CRICKETS, DARKLING BEETLES,
GRASSHOPPERS, SOWBUGS AND ANTS.
ALSO FOR THE CONTROL OF MOLE CRICKETS.

CONTAINS SEVIN® BRAND CARBARYL INSECTICIDE

ACTIVE INGREDIENT:		
Carbaryl: 1-Naphthyl N-methylcarbamate	5.0%	
INERT INGREDIENTS:	95.0%	
	TOTAL	100.0%

*Sevin® is a registered trademark of Rhone-Poulenc for its brand of carbaryl insecticide.

**KEEP OUT OF REACH OF CHILDREN
AND DOMESTIC ANIMALS**

CAUTION

STATEMENTS OF PRACTICAL TREATMENTS
IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention. **IF ON SKIN:** Remove contaminated clothing. Wash affected area with soap and water.
NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
CAUTION: HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust. Do not take internally. Avoid contact with skin or eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. **AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS, AND PONDS DURING APPLICATION OR WHEN CLEANING EQUIPMENT.**
ENVIRONMENTAL HAZARDS
 This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

EPA EST. NO. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. corresponds to first digit in Lot No. on side of label.

Southern Agricultural Insecticides, Inc.

Palmtoe, FL 34220
EPA Reg. No. 829-285


Hendersonville, NC 28753
J1

Beems, NC 28607
NET WEIGHT: 4 LBS.

ACCEPTED

JUN - 2 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 829-285



BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply granules as a broadcast application over the soil with ground equipment. Treatments may be repeated as necessary. Dosages from 1/2 lb. to 1 lb. per 1,000 sq. ft. will control Ants, Cutworms, Armyworms, Crickets, Darkling Beetles, Grasshoppers and Sowbugs. "Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels."

CROP	Dosage Rate 1,000 Sq. Ft.	Days Between Final Application & Harvest
Cucumbers, Squash, Melons, Peanuts	1/2 lb.	None
*Alfalfa, *Peas, & **Cotton	3/4 lb.	*3 Days **7 Days
Beans, Carrots, Corn, Eggplant, Okra, Peppers, Potatoes, Sweet Corn, Tomatoes	1 lb.	None
Asparagus, Strawberries	1 lb.	1 Day
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce, Garden Beets (roots), Horseradish, Parsnips, Rutabagas, Radishes, and Turnips (roots)	1 lb.	3 Days
Sugar Beets, Collards, Endive, Garden Beets (tops), Kale, Leaf Lettuce, Parsley, Spinach, Swiss Chard, Turnips (tops)	1 lb.	14 Days
Tobacco	1/2 - 1 lb.	None
Fruits — Apples, Apricots, Cherries, Citrus, Grapes, Nectarines, Peaches, Pears, Plums Ornamentals — Annual plants, herbaceous plants and shrubs, trees, woody plants	1 lb.	1 Day: Apples, Cherries, Peaches, Pears, Plums 3 Days: Apricots, Nec- tarines; 5 Days: Citrus

LAWN, TURF AND ORNAMENTAL PLANTS

Treatments may be repeated as necessary to control Cutworms, Armyworms, Field Crickets, Mole Crickets, Darkling Beetles, Grasshoppers and Sowbugs.

LAWNS AND TURF — Treat entire area evenly at a dosage of 3/4 to 1 lb. per 1000 Sq. Ft. Apply with a rotary spreader to obtain uniform coverage, giving particular attention to edges of sidewalks and along porch and patio foundations. For best results cut lawn, remove clippings and water thoroughly before treatment.

ORNAMENTAL PLANTINGS — Apply evenly to soil around plants, shrubs and trees at the rate of 3/4 to 1 lb./1000 Sq. Ft. For best results, areas to be treated should be damp.


IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug, and Cosmetic Act permit the sale of crops bearing probable Sevin residues when SEVIN is used in accordance with label directions. If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage, and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk, or meat in excess of established tolerances.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in a cool, dry, locked place out of reach of children and domestic animals. **DISPOSAL:** Do not reuse empty container. Wrap and put in trash.

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.



0 51538 02974 2

BEST AVAILABLE COPY

