

OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

April 9, 2025

Kasey Ray Registrations Southern Agricultural Insecticides P.O. Box 218 Palmetto, FL 34220

Subject: Registration Review Label Amendments Incorporating Mitigation from the

National Marine Fisheries Services (NMFS) Biological Opinions on the Effects of

Carbaryl

Product Name: Cutworm & Cricket Bait EPA Registration Number: 829-285

Application Date: 4/25/24 Decision Number: 595707

Dear Kasey Ray:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application of the above referenced product in connection with the NMFS' Biological Opinion (BiOp) on the effects of carbaryl on federally listed endangered and threatened species and designated critical habitat in accordance with section 7 of the Endangered Species Act (ESA) of 1973, as amended (16 U.S.C. 1531 et seq.). The Agency has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or

Page 2 of 2 EPA Reg. No. 829-285 Decision No. 595707

claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

The Agency received the U.S. Fish and Wildlife Service's (FWS) BiOp on the effects of EPA's national registration of carbaryl and posted it on the <u>website</u> on April 9, 2025. The Agency plans to incorporate and implement this FWS BiOp when issuing the interim decision for carbaryl. EPA will notify the registrants of any need to submit label amendments within 60 days of the issuance of the interim decision.

If you have any questions about this letter, please contact Carolyn Smith at smith.carolyn@epa.gov.

Sincerely,

Jackyn Pyrie

Jaclyn Pyne, Team Leader Risk Management and Implementation Branch 3 Pesticide Re-Evaluation Division Office of Pesticide Programs

Enclosure



MASTER LABEL PART A-INTENDED FOR COMMERCIAL USE

CARBARYL

GROUP

INSECTICIDE

CUTWORM & CRICKET BA

FOR THE CONTROL OF MOLE CRICKETS, CUTWORMS, ARMYWORMS, FIELD CRICKETS, DARKLING BEETLES, GRASSHOPPERS AND SOWBUGS IN COMMERCIAL TURF, RESIDENTIAL LAWNS, ORNAMENTALS and VEGE-TABLES.

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate)	5.00%
INERT INGREDIENTS	95.00%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS **CAUTION**

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN or CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment advice. You may also contact 1-800-858-7378 for emergency medical treatment information.

NOTE TO PHYSICIAN: Carbaryl is an N-methyl carbamate which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Harmful is absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are Barrier Laminate, Butyl Rubber, Neoprene Rubber, Poy-vinyl Chloride (PVC) or Viton. If you want more options, follow the instructions for category A on an EPA chemical-resis-tance category selection chart. Loaders, applicators and other handlers must wear: long-sleeved shirt and long pants, chemical resistant gloves, and shoes plus socks.

Aerial application is prohibited..

ACCEPTED

04/09/2025

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 829-285

Southern Agricultural Insecticides, Inc.

Palmetto, FL. 34220

Hendersonville, N.C. 28793

Boone, N.C. 28607

EPA Reg. NO. 829-285

Net Weight: 40lb. (18.14 kg.)

USER SAFETY REQUIREMENTS

Follow manufacturers' instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When applicators use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(5), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 48 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from area treated. Drift and runoff may kill aquatic invertebrates in water adjacent to treated areas. Do not contaminate water by cleaning equipment or disposing of wastes. Do not contaminate water when disposing of equipment washwater or rinsate. BEE CAUTION: MAY KILL HONEYBEES AND OTHER BEES. This product is highly toxic to honeybees and other bees. Do not apply to plants in bloom.

DIRECTIONS FOR USE

Do not apply this product for agricultural uses in Hawaii.

It is a violation of Federal Law to use this product in a manner Inconsistent with its labeling. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

This product may be applied only using a push spreader. This product may not be applied by hand, spoon, or belly grinder. Product must be watered in immediately following application to lawns. For outdoor use only. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not allow people or pets to enter or allow others allow people or pets to enter the treated area until dusts have settled. Watering-in is required after application to lawns. Do not enter or allow others to enter treated lawns (except those persons involved in the application and watering) until the watering in is complete and until the surface is dry after the watering-in. Do not allow the product to enter any drain during or after application. Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Do not apply or irrigate to the point of runoff. Do not apply to vertical surfaces directly above pervious or impervious surfaces that drain into ditches, storm drains, gutters, or surface waters. Do not apply during rain. Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams or ephemeral streams when water is present, wetlands or natural ponds, estuaries, and commercial fish farm ponds).

All outdoor applications, unless permitted elsewhere on this label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: 1.Applications to soil, lawn, turf, or vegetation. 2. Perimeter band treatments of 6 ft wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn); 3. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch; 4. Applications to vertical surfaces directly above pervious surfaces, such as bare soil, lawn,

turf, mulch or other vegetation, that do not drain into ditches, storm drains, gutters, or surface waters. Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft).

ENDANGERED AND THREATENED SPECIES PROTECTION REQUIREMENTS: Before using this product, you must obtain any applicable Endangered Species Protection Bulletins ('Bulletins') within six months prior to or on the day of application. To obtain Bulletins, go to Bulletins Live! Two (BLT) at https://www.epa.gov/pesticides/bulletins. When using this product, you must follow all directions and restrictions contained in any applicable Bulletin(s) for the area where you are applying the product, including any restrictions on application timing if applicable. It is a violation of Federal law to use this product in a manner inconsistent with its labeling, including this labeling instruction to follow all directions and restrictions contained in any applicable Bulletin(s). For general questions or technical help, call 1-844-447-3813, or email ESPP@epa.gov

REPORTING ECOLOGICAL INCIDENTS: For guidance on reporting ecological incidents, including death, injury, or harm to plants and animals, including bees and other non-target insects, see EPA's Pesticide Incident Reporting website: https://www.epa.gov/pesticide-incidents or call (941)722-3285.

Resistance-management recommendations: For resistance management, **Cutworm Cricket Bait** contains a Group 1A insecticide. Any insect population may contain individuals naturally resistant to **Cutworm Cricket Bait** and other Group 1A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry intervals. The requirements in this box only apply to uses of this product; that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours unless otherwise specified in the directions for use associated with each crop. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: 1. Coveralls over short-sleeve shirt and short pants, 2. Chemical-resistant footwear, shoes plus socks, 3. Chemical-resistant gloves made of any waterproof material, 4. Chemical-resistant headgear for overhead exposure.

When the Restricted-Entry Interval for a crop is 7 days or longer, you must notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, nurseries, or greenhouses. Do not enter or allow others to enter the treated area until dusts have settled. In addition if directions for use require watering-in, do not enter or allow others to enter the treated area (except those involved in the watering-in) until the watering-in is completed and the area has dried.

This product may be applied only by a commercial ground spreader. Product must be watered in immediately after application to residential lawns. Do not apply by hand, spoon, or belly grinder. For outdoor use only.

TO CONTROL MOLE CRICKETS IN TURF, ORNAMENTALS and VEGETABLES

TURF -Including golf courses, cemeteries, commercial landscape, residential lawns and sod farms (REI for sod farms is 24hours.) Treat entire area evenly and uniformly using a commercial type spreader at a dosage of .75 to .9 lb. per 1000 sq. ft. (32 to 40 lbs per acre) Do not make more than 4 applications per year. Do not retreat before 7 days. For best results cut turfgrass, remove clippings and water thoroughly before treatment. When product is applied to residential lawns, watering in immediately after application is required.

ORNAMENTAL PLANTINGS - Apply evenly and uniformly to soil around flowers, shrubs and trees at the rate of .75 to .9 lb./1000 sq. ft. Do not make more than 2 applications per year. Do not retreat before 7 days. For best results, areas to be treated should be damp. (REI is 12 hours. EXCEPTION: the REI is 18 days for ornamentals grown for cuttings (cut flowers or cut foliage) where production is in outdoor areas and where average annual rainfall is less than 25 inches a year.) Notify workers of the application and restricted-entry interval by warning them orally and by posting warning signs at entrances to treated area.. Note: information of average annual rainfall for your area is available from any near-by weather bureau, such as one located at a local airport or one affiliated with the National Oceano-graphic and Atmospheric Administration (NOAA)

VEGETABLE AND OTHER CROPS- Apply granules as a broadcast application over the soil with commercial type ground spreader. Dosages from .45 lb. to .9 lb. per 1,000 sq. ft. (20 to 40 lbs per acre) will control Ants (except fire ants, pharoah ants, harvester ants and carpenter ants), Cutworms, Armyworms, Crickets, Darkling Beetles, Grasshoppers and Sowbugs.

VEGETABLE AND OTHER CROPS	1,000 Sq. Ft.	1 Acre	Max Applications/ Year
Cucumbers (3), Squash (3), Melons (3), *7,**6	.45 lb.	20 lbs.	4
Eggplant (3), Peppers (3), Tomatoes (3) *7, **4	.9 lb.	40 lbs.	4
Broccoli (3), Brussels Sprouts (3), Cabbage (3), Cauliflower (3), *7 **3 Application is permitted only within 30 days from the date of crop emergence or the date of transplanting.	.9 lb.	40 lbs.	3
Garden Beets (roots) (7), Potatoes (7), Radishes and Turnips (roots) (7) Carrots (7), *2, **3	.9 lb.	40 lbs.	3

Number in () behind crop name is the days required between final application and harvest (PHI)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Pesticide should be stored in the original container in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE: To the extent consistent with applicable law, buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

FOR INFORMATION AND QUESTIONS CONCERNING THIS PRODUCT PLEASE CONTACT US BY EMAIL AT info@southernag.com

^{*} behind crop name is Minimum Retreatment Interval, days

^{**} behind crop name is Maximum Number of applications per year

FOR THE CONTROL OF MOLE CRICKETS, CUTWORMS, ARMYWORMS, FIELD CRICKETS, DARKLING BEETLES, GRASSHOPPERS AND SOWBUGS IN LAWNS, ORNAMENTALS, VEGETABLES AND OTHER LISTED CROPS. CONTAINS CARBARYL INSECTICIDE

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl N-methylcarbamate)	5.00%
INERT INGREDIENTS	95.00%
TOTAL	100 00%

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS CAUTION

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN or CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment advice. You may also contact 1-800-858-7378 for emergency medical treatment information.

NOTE TO PHYSICIAN: Carbaryl is an N-methyl carbamate which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS and DOMESTIC ANIMALS

Caution: Harmful if swallowed. Harmful is absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

USER SAFETY RECOMMENDATIONS

Users should wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 48 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate Do not apply when weather conditions favor drift from area treated. Drift and runoff may kill aquatic invertebrates in water adjacent to treated areas. Do not contaminate water by cleaning equipment or disposing of wastes. **BEE CAUTION:** MAY KILL HONEYBEES AND OTHER BEES. This product is highly toxic to hon-eybees and other bees. Do not apply to plants in bloom.

DIRECTIONS FOR USE

Do not apply this product for agricultural uses in Hawaii.

It is a violation of Federal Law to use this product in a manner Inconsistent with its labeling. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

This product may be applied only using a push spreader. This product may not be applied by hand, spoon, or belly grinder. Product must be watered in immediately following application to lawns. For outdoor use only. Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Do not allow people or pets to enter or allow others allow people or pets to enter the treated area until dusts have settled. Watering-in is required after application to lawns. Do not enter or allow others to enter treated lawns (except those persons involved in the application and watering) until the watering in is complete and until the surface is dry after the watering-in. Do not allow the product to enter any drain during or after application. Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment. Do not apply or irrigate to the point of runoff. Do not apply to vertical surfaces directly above pervious or impervious surfaces that drain into ditches, storm drains, gutters, or surface waters. Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams or ephemeral streams when water is present, wetlands or natural ponds, estuaries, and commercial fish farm ponds).

All outdoor applications, unless permitted elsewhere on this label, must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: 1.Applications to soil, lawn, turf, or vegetation. 2. Perimeter band treatments of 6 ft wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn); 3. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch; 4. Applications to vertical surfaces directly above pervious surfaces, such as bare soil, lawn, turf, mulch or other vegetation, that do not drain into ditches, storm drains, gutters, or surface waters.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

Treat surfaces to ensure thorough coverage but avoid runoff.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft).

REPORTING ECOLOGICAL INCIDENTS: For guidance on reporting ecological incidents, including death, injury, or harm to plants and animals, including bees and other non-target insects, see EPA's Pesticide Incident Reporting website: https://www.epa.gov/pesticide-incidents or call (941-722-3285).

Resistance-management: For resistance management, **Cutworm Cricket Bait** contains a Group **1A** insecticide. Any insect population may contain individuals naturally resistant to **Cutworm Cricket Bait** and other Group **1A** insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

TO CONTROL MOLE CRICKETS IN RESIDENTIAL LAWNS, ORNAMENTALS and VEGETABLES

LAWNS- Cover area evenly at a rate of .75 to .9 lb. per 1000 sq. ft. One pound container will treat 1,300 to 1,100 sq. ft. Do not make more than 4 applications per year. Do not retreat before 7 days. For best results cut grass, remove clippings and water thoroughly before treatment. Product must be watered in immediately following application.

ORNAMENTAL PLANTINGS - Cover soil evenly and uniformly around flowers, shrubs and trees at the rate of .75 to .9 lb./1000 sq. ft. One pound will treat 1,300 to 1,100 sq. ft. Do not make more than 2 applications per year. Do not retreat before 7 days. For best results, areas to be treated should be damp. Product must be watered in immediately following application.

VEGETABLE AND OTHER CROPS - Cover area evenly. Treatments may be repeated as listed below. Dosages from .45 lb. to .9 lb. per 1,000 sq. ft. will control Ants (excluding pharaoh ants fire ants, harvester ants and carpenter ants), Cutworms, Army-worms, Crickets, Darkling Beetles, Grasshoppers and Sowbugs. Remove any clippings and water thoroughly before treatment.

VEGETABLE AND OTHER CROPS	1,000 Sq. Ft.	Max Applications/ Year
Cucumbers (3), Squash (3), Melons (3), *7	.45 lb.	4
Eggplant (3), Peppers (3), Tomatoes (3) *7,	.9 lb.	4
Broccoli (3), Brussels Sprouts (3), Cabbage (3), Cauliflower (3), *7 Application is permitted only within 30 days from the date of crop emergence or the date of transplanting.	.9 lb.	3
Garden Beets (roots) (7), Potatoes (7), Radishes and Turnips (roots) (7) Carrots (7), *2,	.9 lb.	3

Number in () behind crop name is the days required between final application and harvest (PHI)

STORAGE AND DISPOSAL

Nonrefillable container. Do not reuse or refill this container. Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Pesticide should be stored in the original container in a locked storage area.

PESTICIDE DISPOSAL: If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

NOTICE: To the extent consistent with applicable law, buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

For information and questions concerning this product please contact us by email at support@southernag.com.

^{*} behind crop name is Minimum Retreatment Interval, days