

829-287

11/5/2009

1/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms Cynthia Huffman
Southern Agricultural Insecticide, Inc
P.O. BOX 218
Palmetto , FL 34220

NOV 05 2009

Subject: Label Notification(s) for Pesticide Registration Notices 2007-4 and 2008-1

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 and 2008-1 dated September 30, 2009 for:

EPA Registration 829-287 SA-50 Liquid Ornamental and Vegetable Fungicide

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and 2008-1, and finds that the label change(s) requested falls within the scope of PRN 2007-4 and 2008-1. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 829-287	2. EPA Product Manager tony Kish	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) SA-50 Liquid Ornamental and Vegetable Fungicide	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: <input checked="" type="checkbox"/> EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

NOV - 5 2009

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change relative to PR Notice 2007-4 and PR Notice 2008-1. See attached page.

Section - III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container Pt, Qt., Gallon _____	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Cynthia Huffman	Title Registrations	Telephone No. (Include Area Code) 941-722-3285
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Cynthia Huffman</i>	3. Title Registrations	
4. Typed Name Cynthia Huffman	5. Date Sep 29, 2009	

**S A SOUTHERN
5 0 AG.**

Notification of label changes relative to PR Notice 2008-1. This notification is consistent with the guidance on PR Notice 2008-1 and the requirements of EPA's regulations at 40 CFR 156.10, 156.80 and 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the guidance of PR Notice 2008-1 and the requirements of 40 CFR 156.10, 156.80 and 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Cynthia Huffman



Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146 and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146 and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Cynthia Huffman

6/10

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
FIRST AID**

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes. Then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Note to Physician: Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

NOTE TO USER: Wear long sleeve shirt, long pants, gloves and goggles or face shield while mixing and applying this product. Keep unprotected persons and animals out of treatment area until spray has completely dried.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT allow people or pets to enter treated areas until sprays have dried or dusts have settled. DO NOT apply this product in a way that will contact other persons, or pets, either directly or through drift. DO NOT combine this product in the spray tank with other pesticides, surfactants, or fertilizers unless your prior use has shown the combination physically compatible, effective and non injurious under your conditions of use. The use of spray stickers or other spray adjuvants is not recommended with this product.

Before opening, slowly invert this container several times to assure uniform mixture. Slowly add the required amount of this product to the spray tank while filling. Agitate and continue to agitate while using.

For handy reference, 1 oz liquid = 2 Tablespoonfuls (TBS) = 6 teaspoonfuls (tsp)

APPLICATIONS INSTRUCTIONS

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (ie., elementary, middle and high schools), campgrounds, churches, and theme parks.

ORNAMENTAL PLANTS:

Apply this fungicide at the rate of 2 1/2 Tablespoonfuls (2 Tablespoonfuls for roses) per gallon of water to cover 350 sq. ft. Apply the spray to run-off when conditions are favorable for disease development. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable. During periods when conditions favor severe disease incidence, generally cloudy wet weather, apply at one week intervals. Apply to plants when foliage and flowers are dry or nearly dry.

Use of this product is recommended for the control of fungal diseases referred to by numbers in parentheses following each ornamental. Ornamentals listed on this label have been tested and found to tolerate applications of chlorothalonil (the active ingredient in this product) at the recommended rates. The user should test for possible phytotoxic responses, using recommended rates on ornamental plants on a small area prior to extensive use. Applications to bloom may damage flowers and/or fruits.

FRUITS AND OTHER STRUCTURES WHICH ARE BORNE ON TREATED PLANTS MUST NOT BE EATEN unless directions for use on the specific food plant are followed as given in the vegetable and fruit sections of this label. Do not feed treated plant parts to livestock. Do not apply more than 27 applications per growing season (year).

ORNAMENTALS RECOMMENDED FOR TREATMENT WITH THIS FLOWABLE FUNGICIDE.

BROADLEAF SHRUBS AND TREES

- Andromeda(Pieris)(4), Ash(Fraxinus)(1), Aspen(1), Azalea(1,2,4), Buckeye(horsechestnut)(1), Cherry-Laurel(1), Crabapple(1,6,8), Dogwood(1), Eucalyptus(3), Euonymus(1), Firethorn(Pyraecantha)(1), Flowering Almond(1,2), Flowering Cherry(1,2), Flowering Peach(1,2), Flowering Plum(1,2), Flowering Quince(1,2), Hawthorn(1,6), Holly (1), Lilac(5), Magnolia, Maple(1), Mountain Laurel(1), Oak(Red group only)(1,7), Oregon-Grape(Mahonia)(6), Photinia(1), Poplar(1), Privet(ligustrum)(1), Rhododendron(1,2,4), Sand Cherry(1,2), Sequoia(1), Spiraea(1), Sycamore, Planetree(1), Viburnum(5), Walnut(Juglans)(1)

FLOWERING PLANTS AND BULBS

(Avoid applications during bloom period on plants where flower injury is unacceptable.)

Arabian Violet (2), Begonia (1), Camellia (2), Carnation (1,2), Chrysanthemum (1,2), Crocus (1), Daffodil (1), Daisy (1), Geranium (1,6), Gladiolus (1,2), Hollyhock (6), Hydrangea (Foliage only)(1,6), Iris (1,2), Lily (1), Marigold (1), Narcissus (1), Pansy (1), Petunia (1,4), Phlox (1), Poinsettia (1)*Discontinue applications to poinsettia prior to bract formation; phytotoxicity is possible on the bracts, Rose (1), Statice (1), Tulip (1), Zinnia (1,5) Do not apply more than 30 applications per season (year).

FOLIAGE PLANTS

Aglaonema(1), Areca palm(1), Artemesia(1), Boston fern(1), Dumbcane(Diffenbachia)(1), Dracaena(1), Fatsia(Aralia)(1), Ficus(1), Florida Ruffle Fern(1), Leatherleaf Fern(1), Lipstick plant(1), Ming aralia(1), Oyster plant(Rhoeo)(1), Parlorpalm, Chamaedorea(1), Peperomia(1), Philodendron(1,4), Prayer plant(Marganta)(1), Syngonium(1), Zebra plant(Aphelandra)(1).

DISEASES CONTROLLED WITH THIS FLOWABLE FUNGICIDE

(1) LEAFSPOTS/FOLIAR BLIGHTS

Actinopelte leafspot	Alternaria leafspot/leaf blight	Anthracoise leaf blotch,spot
Anthracoise(Discula)blight	Ascochyta blight	Bipolaris(Helminthosporium leaf spot)
Botrytis leafspot, leaf blight	Cephalosporium leaf spot	Cercospora leaf spot
Cercosporidium leafspot	Coryneum blight (shothole)	Corynespora leaf spot
Curvularia leaf spot	Cylindrosporium leafspot	Dactylaria leafspot
Didymellina leafspot	Drechslera leafspot	Fabraea(Entomosporium)leafspot
Fusarium leafspot	Gleosporium black leafspot	Ink spot (Drechslera)
Monillinia blossom blight	twig blight	Mycosphaerella ray blight
Myrothecium leafspo,	brown rot	Nematostoma leaf blight
Phyllosticta leaf spot	Rhizoctonia web blight	Ramularia leaf spot
Septoria leaf spot	Sphaeropsis leafspot	Stagonospora leaf scorch
Tan leafspot(Curvularia)	Volutella leaf blight	

(2) FLOWER SPOTS/BLIGHTS

Botrytis flower spot,	flower blight	Curvularia flower spot	Monillinia blossom blight
Ovulinia flower blight	Rhizopus blossom blight	Sclerotinia flower blight	

(3) CYLINDROCLADIUM STEM CANKER

(4) PHYTOPHTHORA LEAF BLIGHT, dieback

(5) POWDERY MILDEWS

Erysiphe cichoracearum	Microsphaera spp
------------------------	------------------

(6) RUSTS

Gymnosporangium spp.	Puccinia spp.	Pucciniastrum hydrangeae
----------------------	---------------	--------------------------

(7) TAPHRINA blister

(8) SCAB (Venturia inaequalis)

THE FOLLOWING ORNAMENTAL PLANT SPECIES WHICH HAVE BEEN TESTED WITH CHLOROTHALONIL AT RECOMMENDED RATES DID NOT EXHIBIT PHYTOTOXICITY.

<u>BOTANICAL NAME</u>	<u>COMMON NAME</u>	<u>BOTANICAL NAME</u>	<u>COMMON NAME</u>
Aechmea fasciata	Aechmea	Episcia cupreata	Flame Violet
Araucaria heterophylla	Norfolk Island Pine	Fittonia spp.	Silver-Nerve Plant
Asplenium nidus	Birdnest fern	Gerbera jamesonii	Gerbera Daisy
Bougainvillea spp.	Bougainvillea	Gynura sarmentosa	Purple Passion Vine
Caladium spp.	Caladium	Gypsophila paniculata	Baby's Breath
Calathea makoyana	Peacock plant	Hoya spp.	Wax Plant
Calistephus chinensis	Aster	Ilex cornuta	Chinese Holly
Carissa grandiflora	Natal plum	Ilex crenata	Japanese Holly
Clerodendron thomsonae	Bleeding heart	Impatiens spp.	Impatiens
Codiaeum spp.	Croton	Pilea cadierei	Aluminum Plant
Cordyline terminalis	TiPlant	Platycerium spp.	Staghorn fern
Crassula argentea	Jade Plant	Sansevieria trifasciata	Birdnest Sansevieria
Cyrtomium falcatum	Holly Leaf Fern	Tolmeia menziesii	Piggy-back Plant
Dionaea muscipula	Venus Fly Trap	Yucca elephantipes	Spineless Yucca
Dizygotheca elegantissima	False Aralia	Zygocactus truncatus	Christmas cactus
Epipremnum aureum	Golden Pothos,	Scindapsus	

NOTE: DO NOT apply this product to either green or variegated Pittosporum or to Schefflera, as multiple applications have been demonstrated to cause phytotoxic responses.

ORNAMENTAL CONIFERS:

Apply this fungicide at the rate given below per gallon of water to cover 350 sq. ft. Apply to obtain uniform coverage of tree canopy. Do not allow livestock to graze treated areas. Rates are given in Tablespoonfuls of product per gallon of water. Do not retreat within 7 days. Do not make more than 6 applications per year for rates up to 4 1/2 TBS. Do not make more than 4 applications per year at rates above 4 1/2 TBS.

<u>DISEASES</u>	<u>TBS/GAL</u>	<u>APPLICATION DIRECTIONS</u>
Swiss needlecast	7 1/2 TBS.	Make first application in Spring when new shoot growth is 1/2 to 2 inches long.
Scleroderris canker(pines)	2 1/2 TBS to 4 1/2 TBS	
Sirococcus tip blight	3 1/2 TBS	
Cyclaneusma and Lophodermium needlecasts	7 1/2 TBS	Apply in early Spring prior to bud break. Repeat applications at 6 to 8 week intervals, until spore release ceases in late Fall. Apply monthly during periods of frequent rainfall. During drought periods applications may be suspended, then resumed upon next occurrence of needle wetness.
Rhabdocline needlecast	2 1/2 TBS to 4 1/2 TBS	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development.
Botrytis seedling blight Phoma twig blight	2 1/2 TBS	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development.

TREE AND ORCHARD CROPS:

Dosages are given in Tablespoonfuls (TBS) per gallon of water. Spray to point of run-off. Obtain complete coverage of the tree canopy.

For stone fruit do not use more than 2.5 gallons of spray per 350 sq. ft. Do not exceed 5 applications per season.

For papaya do not use more than 1 gallon of spray per 350 sq. ft. Do not exceed 3 applications per season. **DO NOT** apply after shuck split and before harvest. If additional fungicide applications are needed before harvest, apply an alternate fungicide labeled for that use.

<u>CROP</u>	<u>DISEASES</u>	<u>TBS/GAL</u>	<u>APPLICATION DIRECTIONS</u>
STONEFRUIT (Peach, Nectarine, Apricot, Cherry, Plum, Prune)	Leaf curl, Coryneum blight (shothole)	2 TBS	Apply at leaf fall in Autumn. Do not retreat within 10 days. When conditions favor disease make one or two additional applications in mid and late Winter before budswell. Where Coryneum blight (shothole) occurs also apply at budbreak and shuck split.
	Brown rot blossom blight	2 TBS	Apply once at popcorn (pink, red, or early white bud) and a second application at full bloom.
	Lacy (russet) scab (plum/prune)		If weather conditions favor disease development make an additional application at petal fall. Do not retreat within 10 days.
	Cherry leaf spot; Peach, Nectarine, Apricot scab	2 TBS	In addition to the bloom applications make one application at shuck split. For control of Cherry leaf spot after harvest, make one application 7 days after fruit has been removed. Where trees have a history of high leaf spot incidence, make a second application 10-14 days later.
Papaya	Alternaria, Fruit rot Anthracnose Stem end rot	3 TBS	Begin treatment when conditions favor disease to 4 TBS development. Do not retreat within 14 days.

VEGETABLE DISEASES:

Rates are given in Tablespoonfuls (TBS) per gallon of water to cover 350 square feet of planting. Spray to obtain thorough coverage but do not exceed 1 gallon of spray mixture per 350 square feet or residue tolerances may be exceeded. The number shown in parentheses following crop name indicates the interval between last application and harvest. When sprays for disease control must be made at intervals closer to harvest than indicated use an alternate fungicide registered for that use. **DO NOT** graze treated areas or feed treated plant parts to livestock.

<u>CROP</u>	<u>DISEASES</u>	<u>TBS/GAL</u>	<u>APPLICATION DIRECTIONS</u>
Bean (snap)(7)	Rust	2 TBS to 4 TBS	Begin applications during early bloom stage or when disease first threatens. Do not retreat within 7 days. Do not exceed 4 applications per season.
	Botrytis blight (gray mold)	4 TBS	
Beans (dry)(42)	Rust	2 TBS	Begin applications during early bloom stage. Do not retreat within 7 days. For use only on beans to be harvested dry with pods removed.
Navy, Pinto, Kidney, Lima, & Blackeye	Anthracnose Downy Mildew Cercospora leaf spot (blackeye only)		Do not exceed 5 applications per season.
Cabbage / Chinese Cabbage(7) (tight headed varieties only)	Alternaria leaf spot	2 1/2 TBS	Begin applications when transplants are set in field, or shortly after emergence for field seeded crop, or when conditions favor disease development. Do not retreat within 7 days. Repeat at 7-10 day intervals.
Cauliflower Broccoli Chinese Broccoli (7) Brussels sprouts	Downy mildew		Do not exceed 8 applications per season.

CROP	DISEASES	TBS/GAL	APPLICATION DIRECTIONS
Carrot (7)	Cercospera (early blight) Alternaria (late blight)	2 1/2 TBS	Start applications when disease threatens. Do not retreat within 7 days. Repeat applications at 7-10 day intervals. Do not exceed 11 applications per season.
Celery(7)	Cercospera (early blight) Septoria (late blight) Basal stalk rot (Rhizoctonia solani)	3 TBS to 4TBS	Weekly applications when transplants are set in field. Do not exceed 8 applications per season. Do not retreat within 7 days.
Corn (sweet)(14)	Helminthosporium leaf blights, Rust	1 TBS to 2 1/2TBS	Begin applications when conditions favor disease development. Do not retreat within 7 days. Repeat at 7 to 10 day intervals. Do not exceed 6 applications per season.
DO NOT APPLY TO SWEET CORN TO BE PROCESSED. DO NOT ALLOW LIVESTOCK TO GRAZE IN TREATED FIELDS. DO NOT ENSILE TREATED CORN OR USE AS LIVESTOCK FORAGE.			
CUCURBITS:	Anthraxnose	2 TBS	Begin applications when plants are in first true leaf stage, or when conditions favor disease development. Do not retreat within 7 days. Do not exceed 7 applications per season.
Cucumber	Downy Mildew	to 2 1/2TBS	
Cantaloupe	Target spot	3 to 4TBS	Under severe conditions, shorten spray intervals.
Muskmelon	Cercospera Leaf spot		
Honeydew	Gummy stem blight		
Watermelon	(black rot)		
Squash	Scab		
Pumpkin	Powdery Mildew (Sphaerotheca only)		
Onion (dry bulb)(7)	Botrytis leaf blight/blast Botrytis neck rot (suppression) Purple blotch Neck rot in storage	1 1/2 to 2 1/2 TBS	Do not retreat within 10 days. Prior to infection shorten spray interval to 7days when disease is present. Do not exceed 11 applications per season. A minimum of 3 applications prior to one week before lifting.
Onion (green bunching)(14)*	Botrytis leaf blight/blast	2 TBS to 4 TBS	Begin weekly applications prior to favorable infection periods and continue for as long as conditions favor disease. Do not retreat within 7 days. DO NOT apply more than 3 times per season to green bunching onions, leeks, or shallots. For Garlic do not exceed 6 applications per season.
Leak (14)*	Purple blotch		
Shallot(14)*	Downey mildew (suppression)		
Garlic (7)			
Parsnip(10)	Alternaria leaf	2 1/2 TBS	Make first application at first sign of disease or when conditions favor disease infection. Do not retreat within 7 days. DO NOT apply more than 4 times per season.
Peanut(14)	Early leaf spot (Cercospera) Late leaf spot (Cercosporidium)	2 TBS	Apply when leaf wetness first occurs or 30-40 days after planting. Do not retreat within 14 days. Do not exceed 8 applications per season.
Potato (7)	Late blight	2 TBS	Rust, Web blotch. Apply as a banded treatment directed over the foliage, beginning when vines are first exposed and leaf wetness occurs. Do not retreat within 5 days. As vines close apply as a complete cover spray at 7-10 day intervals. Use the 7 day interval when disease conditions worsen, being sure to apply 1 gallon for each 350 square feet. Do not exceed 10 applications per season.
Tomato	Early blight, Late blight, Gray Leaf Spot, Septoria leaf spot, Anthracnose, Alternaria fruit rot, Botrytis gray mold, Rhizoctonia fruit rot	3 TBS to 4 TBS	Do not retreat within 7 days. Begin applications when dew or rain occurs and disease threatens. Do not exceed 7 applications per season.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area.
CONTAINER DISPOSAL:
 Nonrefillable container. Do not reuse or refill this container.
 If empty: Do not reuse this container. Place in trash or offer for recycling if available.
 If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

