

CRAIN ABF TOWER AID

ACTIVE INGREDIENT

Foly [oxyethylene (dimethyliminio) ethylene
(dimethyliminio) ethylene dichloride] 15.0%

INERT INGREDIENT 85.0 %

This product contains 1.29 lb. of active ingredient per gallon and weighs 8.58 lb. per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION BEST DOCUMENT AVAILABLE

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin, eyes, or clothing.

FIRST AID: If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Keep out of lakes, streams, or ponds. Permits may be required for discharges containing this pesticide into lakes, streams, ponds or public water. For guidance, contact the regional office of the Environmental Protection Agency.

ACCEPTED

MAY 22 1979

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
EPA UNDER NO.

DIRECTIONS FOR USE
GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CRAIN ABF TOWER AID is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits.

An initial slug addition of 4.0 to 10.0 fluid ounces of **CRAIN ABF TOWER AID** per 1000 gallons of water to provide a concentration of 32 to 80 parts per million of **CRAIN ABF TOWER AID**, based on the total weight of water in the system, is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 1.0 to 10.0 fluid ounces of **CRAIN ABF TOWER AID** per 1000 gallons of water (8 to 80 parts per million of **CRAIN ABF TOWER AID**) should be employed every 2 to 5 days or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

STORAGE & DISPOSAL: Keep container closed when not in use. Do not contaminate water, food, or feed by storage, disposal, or cleaning of equipment. The container must be triple rinsed (or equivalent) and offered for recycling, reconditioning, or disposal in an approved landfill or buried in a safe place. Rinsate that cannot be used or reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

Manufactured by:
CRAIN CHEMICAL COMPANY
2630 Andjori Drive
Dallas, TX 75220

EPA REG. NO. 830-49

EPA EST. No 830 TX-1

NET CONTENTS:

B
A

A
B