

It's Easy to Use

ROBINS COMBINATION FRUIT SPRAY

DIRECTIONS

DOSAGE: Use 5 level tablespoonfuls to 1 gallon of water; 1-pound to 16 gallons of water.

MIXING: Mix the Combination Spray with a small amount of water and stir to a thin smooth paste. Pour into sprayer through a screen. Keep the mixture stirred while spraying.

APPLYING: Apply with any piston type spray pump. Do not use in Rotary pump unless overflow permits full discharge of pump at all time. Cover trees thoroughly and evenly taking special care to wet the top limbs as well as the lower ones. Wet every leaf and fruit.

1 2 APPLES AND PEARS

Codling Moth, Apple Maggot, Green Apple Aphid, Plum Curculio, Red Banded Leafroller, Pear Psylla, Rosy Apple Aphid, usually suppresses European Red and Two Spotted Mites when used in a seasonal program. Scab, Cedar Rust, Quince Rust, Black Rot, Brooks Spot, Fly Speck, Botryosphaeria Rot, Bitter Rot, Sooty Blotch, Apple Blotch. Do not apply within 7 days of harvest.

1. Pre-Bloom (When blossom buds show pink—before any open).
2. As soon as Petals have fallen.
3. Ten days after No. 2.
4. Two weeks after No. 3.
5. Two weeks after No. 4. (Omit this and subsequent applications on early varieties).

Special application for Fall and Winter apples may be advisable 8 or 9 weeks after bloom falls, or 4 weeks after No. 5.

3 4 5 PEACHES, PLUMS AND CHERRIES

Red Banded Leaf Roller, Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Brown Rot, Scab, usually suppresses European Red and Two Spotted Mites when used in a seasonal program. Do not apply within 14 days of harvest on peaches and plums. Do not apply within 30 days of harvest on cherries which are to be dried.

1. As soon as Petals have fallen.
2. Ten days later when fruit shucks are splitting.
3. One week after No. 2.
4. Two weeks after No. 3. (Omit this and subsequent applications on sweet cherry).
5. For late ripening varieties only, such as Elberta and Halberta Giant Peach, make an application 6 weeks after No. 4.

If **CHERRY LEAF SPOT** is a problem (especially in wet summers), spray immediately after harvest and repeat two weeks later if wet weather continues.

YOUNG FRUIT TREES (all kinds listed above) before they start bearing, 2 or 3 applications are sufficient. First application when new growth is about one inch long. Later applications at 2 week intervals.

6 GRAPES

Downy Mildew, Black Rot, Ripe Rot, Omnivorous Leaf Hopper, Grape Leaf Folder.

1. When shoots are 6 to 10" long.
2. 2 weeks later, just before bloom.
3. As soon as blooming is over, or 2 weeks after No. 2.
4. 2 weeks after No. 3.

Do not apply within 14 days of harvest.

This package is sold by weight, and not by volume. When packed it was full as practicable. It may not appear to be full when opened due to settling, but it contains 1-lb.

G. S. ROBINS AND COMPANY
126 Chouteau Avenue
St. Louis, Missouri 63102

Robins COMBINATION Fruit Spray

EPA REG. NO. 842-109

A HOME ORCHARD SPRAY PRODUCT

A combination insecticide and fungicide already mixed—just add the required amount of water and apply as a spray.

ACTIVE INGREDIENTS

N-(mercaptomethyl) phthalimide S-(O, O-dimethyl) phosphorodithioate	12%
Captan (N-(trichloromethyl) thio)-4-cyclohexene-1,2-dicarboximide	12%
Zineb (zinc ethylene bisdithiocarbamate)	6%
INERT INGREDIENTS	70%
Total	100%

CONTENTS: ONE POUND NET . . . MAKES 16 GALS.

WARNING: Keep Out of Reach of Children

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes or on skin, or on clothing. Do not breathe dust or spray mist. In case of contact, wash immediately with soap and water; for eyes, wash copiously with water and get medical attention. Wear clean clothes and wash with soap and hot water before re-use.

Note to Physician: Imidan® is an organophosphorus insecticide. Atropine is antidotal.

Read entire label before using.

Do not contaminate feed and food. Do not graze livestock in treated orchards. Avoid drift to adjoining food and forage crops. Do not apply when weather conditions favor drift from areas treated. Do not re-use empty containers. Destroy by burying in a safe place. Keep this material in cool dry place with top firmly closed. Keep away from fire or live sparks.

Do not apply when fruit or cover crops are in bloom.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

Important Notice—since the use of this product is beyond our control, SELLER makes no warranty of any kind, expressed or implied concerning the use of this product.

Buyer assumes all risk of use or handling, whether in accordance with directions or not.

Use contents of this package within 12 months of time of purchase.

