


**CAUTION: - COMBUSTIBLE MIXTURE**

N. Y. C. Fire Dept. Permit No. X168663



CONTAINS ©BAYGON  
FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

HAUNT INSECT SPRAY is a highly effective residual insecticide formulation for the control of a wide variety of insects, as listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. Haunt Insect Spray drives insects from their hiding places, gives rapid kill and long residual activity.

## INSECT KILLING POWER LASTS FOR WEEKS

FOR SPOT APPLICATION ONLY

## ACTIVE INGREDIENTS

|                                     |        |
|-------------------------------------|--------|
| *o-Isopropoxyphenyl methylcarbamate | 1.00%  |
| Petroleum Distillates               | 83.75% |
| Essential Oils                      | .25%   |

## INERT INGREDIENTS

15.00%

\*U.S. PATENT NO. 3,111,539

THIS MATERIAL USED UNDER LICENSE FROM CHEMAGRO CORPORATION

©BAYGON is the registered trademark of Farbenfabriken Bayer A.G. Chemagro Corporation licensee.

## KILLS

Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Millipedes, Paper Lice, Scorpions, Silverfish, Sowbugs and Spiders. Exposed stages of Caddises, Cigarette Beetles, Dermestids, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Saw-toothed Grain Beetles and Weevils. Use outdoors only for the control of Fleas, Flies, Gnats, Hornets, Mosquitoes, Punkies, Sandflies and Wasps.

USDA REGISTRATION NO. 861-93

SEE DIRECTION PANEL FOR ADDITIONAL PRECAUTIONS  
**CAUTION: KEEP OUT OF REACH OF CHILDREN**

Read Directions Before Use.

CONTENTS 1 GALLON



© 1968

E.I. du Pont de Nemours &amp; Co.

ESTABLISHED 1820

NEW YORK 27, N. Y.

# CAUTION: - COMBUSTIBLE MIXTURE

N.Y.C. FIRE DEPT. PERMIT No. X168663

## DIRECTIONS FOR USE

This is an especially designed residual insecticide for industrial and institutional use only. When used in accordance with the following directions outstanding results are achieved in controlling the named insects in food processing plants, meat processing plants and in research facilities.

**GENERAL:** Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborage. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

**COCKROACHES:** Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables, behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

**ANTS:** Apply to trails and places where ants enter premises such as window and door sills.

**CADELLES, CIGARETTE BEETLES, DERMESTIDS, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES AND WEEVILS (EXPOSED STAGES):** Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

**SILVERFISH, FIREBRATS AND PAPERLICE:** Apply to localized areas infested by these insects.

**BROWN DOG TICKS:** Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

**CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORFIONS AND SPIDERS:** Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention to window sills.

### FOR OUTDOOR USE:

**FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES AND SANDFLIES:** Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

**ANTS, CLOVER MITES, CRICKETS, EARWIGS, HORNETS, MILLIPEDES, SOWBUGS, WASPS AND BROWN DOG TICKS:** Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios, or other places where insects may enter. For ants, spray hills and runways. For control of wasps and hornets apply after dark when all of insects have returned to the nest. Repeat as necessary. Do not spray vegetation.

**FOR USE IN MEAT PACKING AND FOOD PROCESSING PLANTS:** Use only in the inedible product areas of meat packing plants, including those that are federally inspected. Any exposed food products, equipment, or packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

**INSECT CONTROL IN RESEARCH FACILITIES:** The use of Residual Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

### CAUTION

May be harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Avoid breathing spray mist.

Should illness occur, get prompt medical aid.  
To physician - Atropine sulfate is antidotal.

### RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment, and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants. Do not use in edible product areas of food processing plants.

® BAYGON is the registered trademark of Farbenfabriken Bayer A.G. Chemagro Corporation licensee.

**ACCEPTED**  
JUN 15 1969  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER  
ED UNDER NO. 861-93