

861-93

09/04/1974

1/1

Best Available Copy

CAUTION: - COMBUSTIBLE MIXTURE

N. Y. C FIRE DEPARTMENT PERMIT No. X168643

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

This is an especially designed residual insecticide for industrial and institutional use only. When used in accordance with the following directions outstanding results are achieved in controlling the named insects in food processing plants, meat processing plants and in research facilities.



DIRECTIONS FOR USE IN OTHER THAN FOOD PROCESSING ESTABLISHMENTS:

GENERAL: Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages around buildings. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

COCKROACHES: Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables, behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

ANTS: Apply to trails and places where ants enter premises such as around window sills and door frames.

CADELLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, GRANARY WEEVILS and RICE WEEVILS: To aid in the control of and kill the exposed stages of these insects thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.
SILVERFISH, FIREBRATS and PAPER LICE: Apply to localized areas infested by these insects.
BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS and SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention around window sills.

FOR OUTDOOR USE:

PLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES and SANDFLIES: Spray infested areas thoroughly including outside surfaces of screens, doors, ~~and~~ around window frames, foundations, patios and other areas where insects may enter.

ANTS, CLOVER MITES, CRICKETS, EARWIGS, HORNETS, MILLIPEDES, SOWBUGS, WASPS and BROWN DOG TICKS: Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios or other places where insects may enter. For Ants, spray hills and runways. For control of Wasps and Hornets spray after dark when all of the insects have returned to their nest. Repeat as necessary. Do not spray vegetation.

INSECT CONTROL IN RESEARCH FACILITIES: The use of this Residual Insect Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

DIRECTIONS FOR USE WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS:

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing) preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation would also be considered a food area).

Prior to application de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switches in the ON position.

Apply a small amount of this residual insecticide, with equipment capable of delivering a pin thin stream of this material, directly into cracks and crevices, expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housings, junction boxes or switch boxes, conduits or hollow equipment legs where Ants, Centipedes, Cockroaches, Clover Mites, Firebrats, Millipedes, Silverfish, Sowbugs and Spiders hide. Care should be taken to avoid depositing the product on to exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS INSECTICIDE IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS. OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED.

NON-FOOD AREAS - Includes garbage rooms, lavatories, floor drains (to sewers entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Apply to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pillars and similar areas where insects hide or through which they may enter.


DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms and face, thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Avoid breathing spray mist.

If poisoning occurs, get prompt medical attention. To physician - Atropine sulfate is antidote.

This product is toxic to birds and other wildlife. Keep out of any body of water. Do not apply where surface is likely to occur.



RESTRICTIONS
Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment and food handling surfaces. Do not apply to animals.