
861-93



1/2

OCT 7 1974

UNDER THE FEDERAL MONTHCIDE FUNGICIDE AND MODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ED UNDER NO. ______ SUBJECT TO ATTACHED COMMENTS.

2



N. Y. C. FIRE DEPARTMENT PERMIT No. X168663 DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME



CONTAINS ® BAYGON FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY

HAUNT_INSECT SPRAY is a highly effective residual insecticide formulation for the control of a wide variety of insects, as listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. Haunt Insect Spray drives insects from their hiding places, gives rapid kill and long residual activity.

INSECT KILLING POWER LASTS FOR WEEKS

FOR SPOT APPLICATION ONLY

ACTIVE INGREDIENTS

*o-Isopropoxyphenyl methylcarbamate 1.00% Petroleum Distillates 83.75% Essential Oils .25% INERT INGREDIENTS 15.00%


*U.S. PATENT NO. 3,111,539

THIS MATERIAL USED UNDER LICENSE FROM CHEMAGRO CORPORATION

[®] **BAYGON** is the registered trademark of Farbenfabriken Bayer A.G. Chemagro Corporation licensee.



Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Millipedes, Paper Lice, Scorpions, Silverfish, Sowbugs and Spiders. Aids in the control of and kills the exposed stages of Cadelles, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils and Rice Weevils. Use outdoors only for the control of Fleas, Flies, Gnats, Hornets, Mosquitoes, Punkies, Sandflies and Wasps.



CAUTION: -- COMBUSTIBLE MIXTURE I. Y. C. FIRE DEPARTMENT PERMIT No. X168663

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

This is an especially designed residual insecticide for industrial and institutional use only. When used in accordance with the following directions outstanding results are achieved in controlling the named insects in food processing plants, meat processing plants and in research facilities. DIRECTIONS FOR USE IN OTHER THAN FOOD PROCESSING ESTABLISHMENTS. GENERAL: Apply with

a low-pressure sprayer equipped with a coarse spray nazzle. Spray surfaces only until wet, Avoid excessive runoff. Apply to all insect breading sites and harborages around buildings. Also treat adjacent exposed sur-faces where insects may crowl when they come out of hiding. Repeat treatment as necessary. COCKROACHES: Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables, behind and baneath cabinets, refrig-erators, sinks and staves; in and around waste containers, switch panels, junction baxes, floor drains and other places where cockroaches hide or rest.

ANTS: Apply to trails and places where ants enter premises such as around window frames and door frames. CADELES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, GRANARY WEEVILS and RICE WEEVILS: To aid in the control of and kill the exposed stages of these insects thoroughly treat floors and

the walls of the infested area with attention to cracks, crevices, correst and perimeters. Sanitation and dust control are of primary importance. Funigate or destroy infested products. SILVERFISH, FIREBRATS and PAPER LICE: Apply to localized areas infested by these insects. BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after surviva. Do not survey animals. ing. Do not spray animals. \$ pray

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS and SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention, around window frames.

FOR OUTDOOR USE: FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES AND SANDFLIES: Spray infested areas thoroughly includ-ing outside surfaces of screens, doors, around window frames, foundations, patios and other areas where insects mov enter.

ANTS, CLOVER MITES, CRICKETS, EARWIGS, HORNETS, MILLIPEDES, SOWBUGS, WASPS and BROWN DOG TICKS: Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios or other places where insects may enter. For Ants spray hills and runways. For control of Wasps and Hornets spray after dark when all of the insects have returned to their nest. Repeat as necess-Do not spray vegetation.

INSECT CONTROL IN RESEARCH FACILITIES: The use of this Residual Insect Spray in and around re-search animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

DIRECTIONS FOR USE WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS - FOOD HAND-ING ESTABLISHMENTS - Places other than private residences in which exposed food is held, processed,

prepared or served. FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT

ONLY. Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing) preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation would also be considered a food area).

Prior to application de-energize electrical equipment. After application, allow time for oil to evaporate before

÷

starting motors or placing switches in the ON position. Apply a small amount of this Residual Insect Spray, with equipment capable of delivering a pin-thin stream of this material, directly into cracks and crevices, expansion joints between different elements of construc-tion or between equipment bases and the floor, wall voids, motor housings, junction boxes or switch boxes, conduits or hollow equipment legs where Ants, Centipedes, Cockroaches, Clover Mites, Firebrats, Milli-pedes, Silverfish, Sawbugs, and Spiders hide. Care should be taken to avoid depositing the product on to exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing

APPLICATIONS OF THIS INSECTICIDE IN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED.

NON-FOOD AREAS. Includes garbage rooms, lavatories, floor drains (to sewers entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Apply to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter. DO NOT REUSE CONTAINER - DESTROY WHEN EMPTY

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Do not get in eyes, on skin or on clothing, In cose of contact, wash immediately with soap and warm water. Wash all hands, arms and face tharoughly with soap and water before eating or smoking. Wash all contaminated clathing with soap and hat water before -use. Avoid breathing spray mist.

If poisoning occurs, get prompt medical attention. To physician - Atropine sulfate is antidotal. RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contamination of feed, food, food processing equipment and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants.

This product is toxic to birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. (BAYGON is the registered trademork of Farbenfabriken Bayer A. G. Chemagro Corporation licensee. B705-B716

VCCELLEU,