


CAUTION: - COMBUSTIBLE MIXTURE

N. Y. C. FIRE DEPARTMENT PERMIT No. X168663

DO NOT USE. POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME



CONTAINS ©BAYGON

WITH THE ADDITION OF ORTHODICHLOROBENZENE FOR DEODORIZING

A highly effective residual insecticide and deodorant formulation for the control of a wide variety of insects, as listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. This combination deodorant and insect spray drives insects from their hiding places, gives rapid kill and long residual activity.

INSECT KILLING POWER LASTS TWO WEEKS

FOR SPOT APPLICATION ONLY

ACTIVE INGREDIENTS

*o-Isopropoxyphenyl methylcarbamate	1.00%
Petroleum Distillates	73.75%
Orthodichlorobenzene	10.00%
Essential Oils	.25%
INERT INGREDIENTS	15.00%

*U.S. PATENT No. 3,111,539

THIS MATERIAL USED UNDER LICENSE FROM CHEMAGRO CORPORATION

©BAYGON is the registered trademark of Farbenfabriken Bayer A.G. Chemagro Corporation licensee.

KILLS

Anic, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Millipedes, Paper Lice, Scorpions, Silverfish, Sowbugs and Spiders. Aids in the control of and kills the exposed stages of Cadelles, Cigarette Beetles, Drugstore Beetles, Indian Meal Moths, Mediterranean Flour Moths, Red Flour Beetles, Sawtoothed Grain Beetles, Granary Weevils and Rice Weevils. Use outdoors only for the control of Fleas, Flies, Gnats, Hornets, Mosquitoes, Punkies, Sandflies and Wasps.


EPA REGISTRATION No. 861-97

EPA Est. 861-NY-1

CAUTION: SEE DIRECTION PANEL FOR ADDITIONAL PRECAUTIONS KEEP OUT OF REACH OF CHILDREN

Read Directions before Use.

NET CONTENTS: ONE GALLON



DO NOT USE POUR SPILL OR STORE NEAR HEAT OR OPEN FLAME

This is an especially designed residual insecticide for industrial and institutional use only. When used in accordance with the following directions outstanding results are achieved in controlling the named insects in food processing plants, meat processing plants and in research facilities.

DIRECTIONS FOR USE IN OTHER THAN FOOD PROCESSING ESTABLISHMENTS.

GENERAL: Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run off. Apply to all insect breeding sites and harborage. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

COCKROACHES: Thoroughly treat the infested area. Spray baseboards, cracks and crevices, around window and door frames, the under surfaces of shelves, drawers and work tables, behind and beneath cabinets, refrigerators, sinks and stoves, in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

ANTS: Apply to trails and places where ants enter premises such as window and door sills.

CADELLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, SAW-TOOTHED GRAIN BEETLES, GRANARY WEEVILS and RICE WEEVILS: To aid in the control of and kill the exposed stages of these insects thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

SILVERFISH, FIREBRATS and PAPER LICE: Apply to localized areas infested by these insects.

BROWN DOG TICKS: Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, SOWBUGS, SCORPIONS and SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention to window sills.

FOR OUTDOOR USE

FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES and SANDFLIES: Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

ANTS, CLOVER MITES, CRICKETS, EARWIGS, HORNETS, MILLIPEDES, SOWBUGS, WASPS and BROWN DOG TICKS: Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios or other places where insects may enter. For Ants, spray hills and runways. For control of Wasps and Hornets spray after dark when all of the insects have returned to their nest. Repeat as necessary. Do not spray vegetation.

INSECT CONTROL IN RESEARCH FACILITIES: The use of this Residual Insect Spray in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

DIRECTIONS FOR USE WITHIN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS**Limited To Crack And Crevice Treatment Only**

Apply a small amount of this residual insecticide, with equipment that can deliver a pencil thin stream of this material, directly into cracks and crevices, expansion joints between different elements of construction or between equipment bases and the floor, wall voids, motor housings, junction boxes or switch boxes, conduits or hollow equipment legs where Ants, Centipedes, Cockroaches, Clover Mites, Firebrats, Millipedes, Silverfish, Sowbugs and Spiders hide. Care should be taken to avoid depositing the product on to exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

Application of this insecticide in the food areas of food handling establishments other than as a crack and crevice treatment, is not permitted.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, on skin or on clothing. In case of contact, wash immediately with plenty of water. Wash hands, arms and face thoroughly with soap and water before eating, drinking, or smoking. Wash all contaminated clothing with soap and hot water before reuse. Avoid breathing spray mist.

Should illness occur, get prompt medical attention.

To physicians: Atropine sulfate is antidotal.

RESTRICTIONS

Do not use as a space spray. Provide adequate ventilation of areas being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Treatments are not to be used on food products. Avoid contamination of feed, food, food processing equipment and food handling surfaces. Do not apply to animals. Remove pets and aquarium fish tanks before spraying. Do not spray plants.

* BAYGON is the registered trademark of E. I. du Pont de Nemours and Company, Inc., a Delaware corporation.

861-177