DIRECTIONS FOR USE

CROP	INSECT CONTROL AID		Ga 100	Gallons per Appl O Gallons of (Sta		e of cation te of oment)
APPLES	Apple Red Bug European Fruit Lecanium Scale Fruit Tree Leaf			2 2 3	Dormant Dormant Dormant Dormant	
	Roller European Red Mite			2	Green Tip to Delayed Dormant % " Green	
	San Jose Sc Scurfy Scale			3	Delayed Do 1/2 " Green Delayed Do 1/3 " Green	
PEARS	Fruit Tree Leaf 3 Dormant					
·	Pear Leaf B Mite	lister		3	Dormant	
	Pear Psylla			2	Late Dorm	ant ———
PECANS	Obscure Scale		<u></u>	3	Dormant	
PEACHES	Cottony Peach Scale		L.	3 Dormant		
PLUMS & PRUNES	European Red Mite European Fruit Lecanium Scale			2 2 2	Dormant to Delayed Dormant ½" Green	
CROP		% or Gallo of 7E pe 100 Gallon of Water	r		ical or icide	Post Bloom
CITRUS Scales: Glover (long), Chaff, Purple, Red, Yellow, Snow, Brown Soft Scale California Red Scale		1.0-1.3 1.3-1.6				March & April
Scales Listed above & Rust Mite, Greasy Spot, & Sooty Mold (Florida only) California Red Scale		1.0-1.3 1.0-1.3 1.3-1.6	1.0-1.3 1.0-1.3		1 lb. Zineb— 75% WP—or— ¼ Pt. Chloro- benzilate 4	
Scales Listed above & Spider Mites, Black Scale, White Fly California Red Scale (do not use in California)		0.5-0.7 0.5-0.7 0.5-0.7 0.65-0.9		¾ Pt. Ett ¼ Pt. Pa 1 Pt. Mal	rathion 4*	Summer (June- July 15)
Scale Insects in Post Bloom for Oil, Greasy Spot, & Sooty Mold (Florida only)		1.0-1.3				Summer (June- July 15)

*Add Zineb or Chlorobenzitate as listed above for Rust Mite

SEE MIXING DIRECTIONS ON OPPOSITE PANEL



SUNSPRAY 7E

USDA REG. 862-8

ACTIVE INGREDIENT:	BY WEIGHT
PARAFFINIC OIL	99%
INERT INGREDIENTS	1%
	100%

*Unsulfonated Residue	92% min.
Grade of Oil	70 Superior/FC-435-66/ Unclassified
Weight per Gallon	

Sun Oil Company, 1608 Walnut St., Philadelphia, Pa. 19103

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. If swallowed, do not induce vomiting. Avoid Breathing of vapors and spray mists. Do not pollute fish and wildlife waters by rinsing or draining of equipment. See other cautions

55 GALLONS NET · U.S. STANDARD

A-5932 Rev. 1 Printed in U.S.A.

Keep oil container tightly closed in storage to prevent entry of water. Spray only when temperature is above 40°F, and there is no danger of freezing.

ON CITRUS: Do not apply when trees are in or near wilting, Oil applied during August and September may affect solids content, retard color. Oil applied after October 1 may increase cold damage to trees. Observe all cautions and limitations on labels of all products used in mixtures. Do not use 7E in combination with or immediately preceding or following applications of Dinitro compounds, sulfurs, Captan, Folget, Dyrene, Karathane, Morestan, or Sevin. Check tank mixtures for chemical and physical compatibility.

MIXING DIRECTIONS


- 1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
- 2. Add other desired pesticides as listed on left panel mixing thoroughly prior to the addition of oil.
- 3. Add oil under agitation when 3/4 full topping off with water to form milky solution.
- 4. Maintain agitation until solution is used.
- 5. In small equipment lacking agitators—stir or shake diluted spray frequently during application.

TIMING THE TREATMENT: See table on other panel for general use. You must determine final timing to fit local growth and climatic conditions.

DO NOT EXCEED MAXIMUM RATES OR APPLY LATER THAN RECOMMENDED.

TERMS

The manufacturer guarantees this material as true to the label declaration of contents and quality. The manufacturer and the vendor make no further representation or warranty and shall not be held responsible for any injury claimed as resulting from the use of this material. Climatic conditions and other factors, including errors in mixing and application, beyond the manufacturer's and the vendor's control, may after results; therefore the buyer accepts and uses this material under the above terms.



ACCEPTED

8-29