

DIRECTIONS FOR USE

CROP	INSECT CONTROL AID	Application Rate 11E Gallons per 100 Gallons of Dilute Spray	Time of Application (Stage of Development)
APPLES	Apple Red Bug	2	Dormant
	European Fruit Lecanium Scale	2	Dormant
	Fruit Tree Leaf Roller	3	Dormant
	European Red Mite	2	Green Tip to Delayed Dormant
	San Jose Scale	2	1/2" Green
PEARS	Fruit Tree Leaf Roller	3	Dormant
	Pear Leaf Blister Mite	3	Dormant
PEACHES	Obscure Scale	3	Dormant
	Cottony Peach Scale	3	Dormant
PLUMS & PRUNES	European Red Mite	2	Dormant to Delayed Dormant
	European Fruit Lecanium Scale	2	1/2" Green
NECTARINES	San Jose Scale	2	Dormant
CHERRIES (sweet & tart)	San Jose Scale	2	Dormant
BLUEBERRIES	San Jose Scale	2	Dormant
	Putnam Scale	3	Dormant
AZALEA	Azalea Bark Scale	2	Dormant to Bud Break
	European Red Mite	2	Dormant to Bud Break
FLOWER CRAB	As Above	2	As Above
FLOWERING CHERRY	As Above	2	As Above
DOGWOOD	Dogwood Scale	2	As Above
	San Jose Scale	2	As Above
EUONYMUS	San Jose Scale Euonymus Scale	2	As Above
HOLLY	Holly Scale	2	As Above
HAWTHORN	San Jose Scale	2	As Above
JUNIPER	Juniper Scale	2	As Above
LILAC	San Jose Scale	2	As Above
PINES	Pine Needle Scale	2	As Above
PRIVET	White Peach Scale	3	As Above
RHODO-DENDRON	Azalea Bark Scale	2	As Above
ROSE	Rose Scale	2	Dormant
SPIREA	San Jose Scale	2	Dormant to Bud Break

SEE MIXING DIRECTIONS ON OPPOSITE PANEL



SUNSPRAY 11E

EPA REG. 862-319

ACTIVE INGREDIENT:	BY WEIGHT
PARAFFINIC OIL	98.8%
INERT INGREDIENTS	1.2%
	100%

Unulfonated Residue 92% min.
Grade of Oil 100 Superior/Unclassified
Weight per Gallon 7.2 lbs.

Sun Oil Company, 1608 Walnut St., Philadelphia, Pa. 19103

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. If swallowed, do not induce vomiting.
Avoid Breathing of vapors and spray mists. Do not pollute fish and wildlife waters by rinsing or draining of equipment. See other cautions

55 GALLONS NET • U.S. STANDARD

A-6086 Printed in U.S.A.

CAUTION

Keep oil container tightly closed in storage to prevent entry of water. Spray only when temperature is above 40°F. and there is no danger of freezing. Store drum on side. Roll drum before use if stored overseason.

GENERAL: Do not apply when plants are in or near wilting. Observe all cautions and limitations on labels of products recommended for mixture with oil.

DO NOT USE 11E in combination with or immediately preceding or following applications of Chloropropylate, Captan, Dinitro Compounds, Dyrene, Dikar, Dichlone, Folpet, Karathane, Morestan, Morocide, Sulfur Compounds, Omite and Sevin. Check tank mixtures for physical and chemical compatibility. Mix only enough material to do the job.

ORNAMENTALS: Do not use on Douglas Fir, Sugar and Japanese Maple, Beech, Birch, Hickory, Walnut or Butternut. Oil will remove blue color from glaucous bearing plants (Blue Spruce).

MIXING DIRECTIONS


1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides mixing thoroughly prior to the addition of oil.
3. Add oil under agitation when 3/4 full topping off with water to form milky solution.
4. Maintain agitation until solution is used.
5. In small equipment lacking agitators—stir or shake diluted spray frequently during application.

TIMING THE TREATMENT: See table on other panel for general use. You must determine final timing to fit local growth and climatic conditions.

DO NOT EXCEED MAXIMUM RATES OR APPLY LATER THAN RECOMMENDED.

TERMS

Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer/user accepts the product on these conditions. Timing and method of applications, weather, and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller.



862-9

862-9

10-18-72

10-1