


55 GALLONS NET
U.S. STANDARD



Superior Spray Oil 6E

USDA REG. 862-

ACTIVE INGREDIENT:	BY VOLUME	BY WEIGHT
Refined Hydrocarbon*	99%	98.9%
Inert Ingredients:	1%	1.1%
	100%	100%

*Unsulfonated Residue.....92% min.
Grade of Oil.....60 Superior/FC-412-66

SUN OIL COMPANY • PHILADELPHIA, PA. 19103

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. See other cautions →

Litho in U.S.A.

DIRECTIONS FOR USE

CROP	INSECTS CONTROLLED	Rate of Application (Gallons/E Spray Oil/ 100 Gallons Dilute Spray)	Time of Application (Stage of Development)
APPLES	Apple Red Bug	2	Dormant
	European Fruit	2	Dormant
	Lecanium Scale	2	Dormant
	Fruit Tree Leaf Roller	3	Dormant
	European Red Mite	2	Green Tip to Delayed Dormant 1/2" Green
	San Jose Scale	2	Delayed Dormant 1/2" Green
	Scurfy Scale	3	Delayed Dormant 1/2" Green
PEARS	Fruit Tree Leaf Roller	3	Dormant
	Pear Leaf Blister Mite	3	Dormant
	Pear Psylla	2	Late Dormant
PEACHES	Cottony Peach Scale	3	Dormant
PLUMS & PRUNES	European Red Mite	2	Dormant to Delayed Dormant 1/2" Green
	European Fruit	2	
	Lecanium Scale	2	
CITRUS	Rust Mite, Spider Mites and Scale	1*	Post Bloom March-April
	Rust Mite, Greasy Spot and Scales	1*	Summer Spray June-July 15
*Add zineb or chlorobenzilate, or ethion at better fruit program recommendation.			
SEE MIXING DIRECTIONS ON OPPOSITE PANEL			

CAUTION

Keep oil container tightly closed in storage to prevent entry of water. Spray only when temperature is above 40°F and there is no danger of freezing.

ON CITRUS: Do not apply when trees are in or near wilting. Oil applied during August and September may affect solids content, retard color. Oil applied after October 1 may increase cold damage to trees. Do not use 6E in combination with or immediately preceding or following applications of Dinitro compounds, sulfurs, Captan, Folpet, Dyrene, Kerathane, Morestan, or Sevin. Check tank mixtures for chemical and physical compatibility.

ON ORNAMENTALS: Do not apply to walnut, beech, hickory, or sugar and Japanese maples. Evergreens tend to be sensitive to oils. Use extreme caution and do not overspray.

TERMS

The manufacturer guarantees this material as true to the label declaration of contents and quality. The manufacturer and the vendor make no further representation or warranty and shall not be held responsible for any injury claimed as resulting from the use of this material. Climatic conditions and other factors, including errors in mixing and application, beyond the manufacturer's and the vendor's control, may alter results; therefore the buyer accepts and uses this material under the above terms.

MIXING DIRECTIONS

1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides mixing thoroughly prior to the addition of oil.
3. Add oil under agitation when 3/4 full topping off with water to form milky solution.
4. Maintain agitation until solution is used.
5. In small equipment lacking agitators—stir or shake diluted spray frequently during application.

TIMING THE TREATMENT: See table on reverse side for general use. You must determine final timing to fit local growth and climatic conditions. DO NOT EXCEED MAXIMUM RATES OR APPLY LATER THAN RECOMMENDED.

862-11

8961/4e/60

11