

DIRECTIONS FOR USE

55 GALLONS NET
U.S. STANDARD

CAUTION



CROP	INSECTS CONTROLLED	Rate of Application (Gallons 6E Spray Oil/ 100 Gallons Dilute Spray)	Time of Application (Stage of Development)
APPLES	Apple Red Bug	2	Dormant
	European Fruit	2	Dormant
	Lecanium Scale	2	Dormant
	Fruit Tree Leaf Roller	3	Dormant
	European Red Mite	2	Green Tip to Delayed Dormant ½" Green
	San Jose Scale	2	Delayed Dormant ½" Green
	Scurfy Scale	3	Delayed Dormant ½" Green
PEARS	Fruit Tree Leaf Roller	3	Dormant
	Pear Leaf Blister Mite	3	Dormant
	Pear Psylla	2	Late Dormant
PEACHES	Cottony Peach Scale	3	Dormant
PLUMS & PRUNES	European Red Mite	2	Dormant to
	European Fruit Lecanium Scale	2	Delayed Dormant ½" Green
	Oil	Chemical or Pesticide	Post Bloom
CITRUS	Scales: Glover (long), Choff, Purple, Red, Yellow, Snow, Brow Soft Scale	1.0-1.3%	March & April
	Scales Listed Above & Rust Mite, Greasy Spot, & Sooty Mold (Florida Only)	1.0-1.3%	1 lb. Zineb—75% WP —or— ¼ Pt. Chlorobenzilate 4
	Scales Listed Above & Spider Mites, Black Scale, White Fly	0.5-0.7%	¼ Pt. Ethion 4*
		0.5-0.7%	¼ Pt. Parathion 4*
		0.5-0.7%	1 Pt. Matathlion 5*
	Scale Insects in Post Bloom for Oil, Greasy Spot, & Sooty Mold (Florida Only)	1.0-1.3%	Summer (June-July 15)

*Add Zineb or Chlorobenzilate as listed above for Rust Mite

SEE MIXING DIRECTIONS ON OPPOSITE PANEL

Superior Spray Oil 6E

USDA REG. 862-11

ACTIVE INGREDIENT:	BY VOLUME	BY WEIGHT
PARAFFINIC OIL	99%	98.9%
INERT INGREDIENTS:	1%	1.1%
	100%	100%

*Unulfonated Residue.....92% min.
Grade of Oil.....60 Superior/FC-412-66

Sun Oil Company, 1608 Walnut St., Philadelphia, Pa., 19103

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. See other cautions ➔

Keep oil container tightly closed in storage to prevent entry of water. Spray only when temperature is above 40°F. and there is no danger of freezing.

ON CITRUS: Do not apply when trees are in or near wilting. Oil applied during August and September may affect solids content, retard color. Oil applied after October 1 may increase cold damage to trees. Observe all cautions and limitations on labels of all products used in mixtures. Do not use 6E in combination with or immediately preceding or following applications of Dinitro compounds, sulfurs, Captan, Folpet, Dyrene, Karathane, Morestan, or Sevin. Check tank mixtures for chemical and physical compatibility.

MIXING DIRECTIONS

1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides as listed on left panel mixing thoroughly prior to the addition of oil.
3. Add oil under agitation when ¾ full topping off with water to form milky solution.
4. Maintain agitation until solution is used.
5. In small equipment lacking agitators—stir or shake diluted spray frequently during application.

TIMING THE TREATMENT: See table on other panel for general use. You must determine final timing to fit local growth and climatic conditions.

DO NOT EXCEED MAXIMUM RATES OR APPLY LATER THAN RECOMMENDED.

TERMS

The manufacturer guarantees this material as true to the label declaration of contents and quality. The manufacturer and the vendor make no further representation or warranty and shall not be held responsible for any injury claimed as resulting from the use of this material. Climatic conditions and other factors, including errors in mixing and application, beyond the manufacturer's and the vendor's control, may alter results; therefore the buyer accepts and uses this material under the above terms.

862-11
04/22/1969