



ACCEPTED

MAR 1 1 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND ROENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 862-11. SUBJECT
 TO ATTACHED COMMENTS.

SUNSPRAY 6E

USDA REG. 862-11

ACTIVE INGREDIENT:	BY WEIGHT
PARAFFINIC OIL	99%
INERT INGREDIENTS:	1%
	100%

*Unulfonated Residue 94% min.
 Grade of Oil 60 Superior/FC-412-66

Sun Oil Company, 1608 Walnut St., Philadelphia, Pa., 19103

CAUTION: KEEP OUT OF REACH OF CHILDREN.

If accidentally ingested Do Not Induce Vomiting
 Harmful if swallowed. See other cautions →

55 GALLONS NET • U.S. STANDARD

Litho in U.S.A.

DIRECTIONS FOR USE

CROP	INSECT CONTROL AND	Rate of Application (Gallons OE Spray Oil/100 Gallons Dilute Spray)	Time of Application (Stage of Development)	
APPLES	Apple Red Bug	2	Dormant	
	European Fruit	2	Dormant	
	Locanium Scale	2	Dormant	
	Fruit Tree Leaf Roller	3	Dormant	
	European Red Mite	2	Green Tip to Delayed Dormant	
	San Jose Scale	2	1/2" Green	
PEARS	Fruit Tree Leaf Roller	3	Dormant	
	Pear Leaf Blister Mite	3	Dormant	
	Pear Psylla	2	Late Dormant	
PECANS	Scurfy Scale	3	Dormant	
PEACHES	Cottony Peach Scale	3	Dormant	
PLUMS & PRUNES	European Red Mite	2	Dormant to	
	European Fruit	2	Delayed Dormant	
	Locanium Scale	2	1/2" Green	
		Oil	Chemical or Pesticide	Post Bloom
CITRUS				
	Scales: Glover (Insg), Chiff, Purple, Red, Yellow, Seaw, Brown Soft Scale	1.0-1.3%		March & April
	California Red Scale	1.3-1.6%		
	Scales Listed Above & Rust Mite, Greasy Spot, & Sooty Mold (Florida Only)	1.0-1.3%	1 lb. Zineb-75% WP	March & April
	California Red Scale	1.0-1.3%	-or- 1/4 Pt. Chlorobenzilate 4	April
	Scales Listed Above & Spider Mites, Black Scale, White Fly	0.5-0.7%	1/2 Pt. Ethion 4*	Summer (June-July 15)
	California Red Scale	0.5-0.7%	1/4 Pt. Parathion 4*	
	Scale Insects in Post Bloom for Oil, Greasy Spot, & Sooty Mold (Florida Only)	0.5-0.9%	1 Pt. Malathion 5*	
		1.0-1.3%		Summer (June-July 15)

*Add Zineb or Chlorobenzilate as listed above for Rust Mite

SEE MIXING DIRECTIONS ON OPPOSITE PANEL

CAUTION

Keep oil container tightly closed in storage to prevent entry of water. Spray only when temperature is above 40°F. and there is no danger of freezing.

ON CITRUS: Do not apply when trees are in or near wilting. Oil applied during August and September may affect solids content, retard color. Oil applied after October 1 may increase cold damage to trees. Observe all cautions and limitations on labels of all products used in mixtures. Do not use 6E in combination with or immediately preceding or following applications of Dinitro compounds, sulfurs, Captan, Folpet, Dyrane, Karathane, Morestan, or Sevin. Check tank mixtures for chemical and physical compatibility.

MIXING DIRECTIONS

1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides as listed on left panel mixing thoroughly prior to the addition of oil.
3. Add oil under agitation when 3/4 full topping off with water to form milky solution.
4. Maintain agitation until solution is used.
5. In small equipment lacking agitators—stir or shake diluted spray frequently during application.

TIMING THE TREATMENT: See table on other panel for general use. You must determine final timing to fit local growth and climatic conditions.

DO NOT EXCEED MAXIMUM RATES OR APPLY LATER THAN RECOMMENDED.

TERMS

The manufacturer guarantees this material as true to the label declaration of contents and quality. The manufacturer and the vendor make no further representation or warranty and shall not be held responsible for any injury claimed as resulting from the use of this material. Climatic conditions and other factors, including errors in mixing and application, beyond the manufacturer's, and the vendor's control, may alter results; therefore the buyer accepts and uses this material under the above terms.

862-11

03/11/1971

1/1