



SUNSPRAY 6E

EPA REG. 862-11

CAUTION: KEEP OUT OF REACH OF CHILDREN
Harmful if swallowed. If swallowed, do not induce vomiting.
Avoid breathing of vapors and spray mists. Do not pollute fish and wildlife waters by rinsing or draining of equipment.

STORAGE AND DISPOSAL

- 1. **PROHIBITIONS.** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- 2. **PESTICIDE DISPOSAL.** Pesticide, spray mixture, or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides buried in a safe place away from water supplies.
- 3. **CONTAINER DISPOSAL**
 - a. Reseal container and offer for reconditioning, or
 - b. Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.
- 4. **GENERAL.** Consult Federal, state or local disposal authorities for approved alternative procedures.

	BY WT.
ACTIVE INGREDIENTS: Refined Petroleum Distillate*	98.8%
INERT INGREDIENTS	1.2%
	100 %

*Unsulfonated Residue of Petroleum Distillate **92% Min.**
 Weight per Gallon **7.1 lbs.**

FOR FRUIT TREES • SHADE TREES • SHRUBS • SUGAR BEETS AND SWEET CORN

TERMS—Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer/user accepts the product on these conditions. Timing and method of applications, weather, and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller.

SUN PETROLEUM PRODUCTS COMPANY • Philadelphia, PA 19103

BULK • U.S. STANDARD

CROP	INSECT CONTROL AID	Application Rate 6E Gallons per 100 Gallons of Dilute Spray	Time of Application (Stage of Development)
APPLES	Apple Red Bug	2	Dormant
	European Fruit Lecanium Scale	2	Dormant
	Fruit Tree Leaf Roller	3	Dormant
	European Red Mite	2	Green Tip to Delayed Dormant ½" Green
	San Jose Scale	2	Delayed Dormant ½" Green
	Scurfy Scale	3	Delayed Dormant ½" Green
PEARS	Fruit Tree Leaf Roller	3	Dormant
	Pear Leaf Blister Mite	3	Dormant
	Pear Psylla	2	Late Dormant
PECANS	Obscure Scale	3	Dormant
PEACHES	Cottony Peach Scale	3	Dormant
PLUMS & PRUNES	European Red Mite	2	Dormant to Delayed Dormant ½" Green
	European Fruit Lecanium Scale	2	Dormant to Delayed Dormant ½" Green

CROP	% or Gallons of 6E per 100 Gallons of Water	Chemical or Pesticide	Post Bloom
CITRUS Scales: Glover (long), Chaff, Red, Yellow, Snow, Soft Scale California Red Scale	1.0-1.3		March & April
	1.3-1.6		
Scales Listed above & Rust Mite, Greasy Spot, & Sooty Mold (Florida only) California Red Scale	1.0-1.3 1.0-1.3 1.3-1.6	1 lb. Zineb—75% WP—or—¼ Pt. Chlorobenzilate 4	March & April
Scales Listed above & Spider Mites, Black Scale, White Fly California Red Scale (do not use in California)	0.5-0.7 0.5-0.7 0.5-0.7 0.65-0.9	¾ Pt. Ethion 4* ¼ Pt. Parathion 4* 1 Pt. Malathion 5*	Summer (June-July 15)
Scale Insects in Post Bloom for Oil, Greasy Spot, & Sooty Mold (Florida only)	1.0-1.3		Summer (June-July 15)

*Add Zineb or Chlorobenzilate as listed above for Rust Mite

SEE MIXING DIRECTIONS ON OPPOSITE PANEL

ACCEPTED
DEC 12 1979
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under
EPA Reg. No. 862-11



SUNSPRAY 6E

EPA REG. 862-11

ACTIVE INGREDIENT:	BY WEIGHT
PARAFFINIC OIL	98.8%
INERT INGREDIENTS	1.2%
	100%

Unulfonated Residue 92% min.
Weight per Gallon 7.1 lbs.

Sun Petroleum Products Company - Philadelphia, Pa. 19103

CAUTION: KEEP OUT OF REACH OF CHILDREN
Harmful if swallowed. If swallowed, do not induce vomiting.
Avoid Breathing of vapors and spray mists. Do not pollute fish and wildlife waters by rinsing or draining of equipment. See other cautions

STORAGE AND DISPOSAL

- Prohibitions.** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
- Pesticide Disposal.** Pesticide spray mixture, or rinse that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.
- Container Disposal.** (a) Reseal container and offer for reconditioning, or (b) triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.
- General.** Consult Federal, state or local disposal authorities for approved alternative procedures.



55 GALLONS NET • U.S. STANDARD

Keep oil container in a cool place to prevent entry of water. Temperature is above freezing.

ON CITRUS: Do not apply to wilting. Oil applied to wilting affect solids content. October 1 may in some areas. Observe all cautions on labels of products used in combination with or immediately after applications of DDT, Folpet, Dyrene, Karathion, and other tank mixtures for citrus.

- Add sufficient water for proper agitation.
- Add other desired pesticides mixing thoroughly.
- Add oil under agitation with water to form an emulsion.
- Maintain agitation during application.
- In small equipment, dilute spray first.

TIMING THE TREATMENT: For general use. Yield fit local growth and conditions.

DO NOT EXCEED DIRECTIONS. APPLY LATELY.

Seller's guarantee covers the label and subject to the product on the date of applications, with other warranties with other products, and other conditions. This product are not to be used for any other purpose.