



A superior horticultural spray oil for insect and mite pest management

Active Ingredient: Paraffinic Oil	98.8%
Inert Ingredient: Emulsifier	1.2%
*Unsulfonated Residue of Paraffinic Oil	92.0% Min
*50% Distillation Point of Paraffinic Oil	414.0°F
*10%-90% Distillation Range of Paraffinic Oil	65.0°F Max
*Flash Point	345°F
Weight per Gallon	7.1 lbs.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**PRECAUTIONARY STATEMENTS**

Hazards to humans & domestic animals  
 Avoid contact with eyes, skin and clothing. Harmful if swallowed.  
 Avoid breathing of spray mists or vapors. Avoid contamination of feed and foodstuffs.

**STATEMENT OF PRACTICAL TREATMENT**  
 • If swallowed: Do not induce vomiting. Call a physician immediately.  
 • If on skin: Wash with soap and water.  
 • If in eyes: Flush with water.  
 • If inhaled: Move to fresh air.

**PERSONAL PROTECTIVE EQUIPMENT**  
**APPLICATORS AND OTHER HANDLERS MUST WEAR:**  
 • Long-sleeved shirt and long pants  
 • Shoes plus socks  
 Follow manufacturer's instructions for maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**  
 Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

**ENVIRONMENTAL HAZARDS**  
 This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

**STORAGE AND DISPOSAL**

- Prohibitions:** Do not contaminate water, soil, or feed by storage or disposal. Open dumping is prohibited.
- Storage:** Store in a cool, dry, locked area out of the reach of children. Keep all containers tightly closed in storage to prevent entry of water.
- Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- Container Disposal:**  
 Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.  
 Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

EPA Registration No. 862-23



**Sun Company, Inc. (R&M)**  
 Ten Penn Center  
 1801 Market Street  
 Philadelphia, PA 19103-1699

**EPA Establishment No. 862-PA-1**  
**Net Contents: 2.5 Gallons**

**BEST AVAILABLE COPY**

PULL HERE TO OPEN  
PRESS TO RESEAL



72704  
EPA

A superior horticultural spray oil for insect and mite pest management.

Active Ingredient: Paraffinic Oil*	98.8%
Inert Ingredient: Emulsifier	1.2%
*Unulfonated Residue of Paraffinic Oil	92.0% Min
*50% Distillation Point of Paraffinic Oil	414.0°F
*10%-90% Distillation Range of Paraffinic Oil	65.0°F Max
*Flash Point	345°F
Weight per Gallon	7.1 lbs.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

**Statement of Practical Treatment**

- If swallowed: Do not induce vomiting. Call a physician immediately.
- If on skin: Wash with soap and water.
- If in eyes: Flush with water.
- If inhaled: Move to fresh air.

See Additional Precautionary Statements Inside Booklet.  
See Directions For Use Inside Booklet.

Fix-a-Form US PAT NO 4,376,008  
AMERICAN AIR FILTERS



**Sun Company, Inc. (R&M)**  
Ten Penn Center  
1801 Market Street  
Philadelphia, PA 19103-1699

EPA Registration No. 862-23

**BEST AVAILABLE COPY**

378

### PRECAUTIONARY STATEMENTS

#### Hazards to humans & domestic animals

Avoid contact with eyes, skin and clothing. Harmful if swallowed. Avoid breathing of spray mists or vapors. Avoid contamination of feed and foodstuffs.

#### Personal Protective Equipment

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should:  
\*Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### Environmental Hazards

This product is toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

### STORAGE AND DISPOSAL

1. **Prohibitions:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.
2. **Storage:** Store in a cool, dry, locked area out of the reach of children. Keep oil container tightly closed in storage to prevent entry of water.
3. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
4. **Container Disposal:** **Metal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. **Plastic:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### USE PRECAUTIONS

Keep oil container tightly closed in storage to prevent entry of water. All horticultural oils interfere with or slow plant transpiration and respiration during the period of evaporation. Phytotoxicity may result if sprayed to plants during periods of prolonged high temperature and high relative humidity. Do not spray to plants under moisture stress.

Do not use this product with dimethoate (Cygon), or fungicides such as captan (Captan), anilazine (Dyrene), folpet (Folpet), dinocap (Karathane), oxythioquinox (Morestan), or any other product containing sulfur. If possible, either keep the spray equipment used for these compounds separate from the equipment used for oil, or make sure that the sprayer is thoroughly cleaned so that no residue from these compounds remain. Do not use with dimethoate or Sevin 50W formulation (carbaryl) on deciduous fruit trees.

### MIXING INSTRUCTIONS

1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides. If wettable powder formulation, mix the water and powder thoroughly so that the powder is totally suspended in the water before the oil is added. If other pesticide to be added is an emulsifiable formulation, do so after the oil and water has been thoroughly mixed.
3. Add oil under agitation when tank is 3/4 full. Top off with water to form milky solution.
4. Maintain agitation until solution is used.
5. In small equipment lacking agitators, stir or shake diluted spray frequently during application.
6. It is important for users to read and follow all directions and restrictions on the labels of the proposed tank mix products.
7. Flush fluid in sprayer hose lines back into tank reservoir if fluid is allowed to stand for more than 20 minutes.

### GENERAL INFORMATION

This product controls adelgids, aphids, lace bugs, leafhoppers, leafminers (larvae), mealybugs, mites, plant bugs, psyllids, sawfly larvae, scales, whitefly and eggs of aphids, mites and certain caterpillars on vegetables, fruits, tree nuts, certain field crops, shrubs, trees, greenhouse plants, ornamental foliage plants and flowers. This product can be applied up to harvest.

### APPLICATION INSTRUCTIONS

The target pest must be completely covered with spray. Oil residue on the plant surface often acts as a feeding and oviposition deterrent. However, the primary target is the pest itself as oil is a contact pesticide.

**DILUTE APPLICATIONS** (greater than 150 gallons spray per acre) in most cases, ensure the best coverage.

**Note:** This pesticide is to be sold only in this original unbroken container.

**CONCENTRATE APPLICATIONS** (usually from 45 to 125 gallons spray per acre) may reduce coverage and effectiveness. Concentrate application includes the use of low volume (from 10 to 100 gallons spray per acre) air-blast or air-carrier sprayers. A concentrate application can provide satisfactory results as long as the spray unit is properly engineered, calibrated and operated. Speed of travel for ground application is extremely important. Tractor speed from 1 M.P.H. to 4 M.P.H. is recommended depending on crop, crop size and target pest.

**AERIAL APPLICATIONS** Use only as an emergency application when soil conditions do not permit regular ground application. Helicopter only. Apply quantity of product shown for each listed crop for control of listed insects in sufficient water to make a minimum of 20 gals. dilute spray per acre.

### TIMING THE TREATMENT

Applicator must determine the precise timing to fit local growth and climatic conditions.

**DO NOT EXCEED MAXIMUM RATES OR APPLY MORE OFTEN THAN RECOMMENDED.**

**MAY BE USED UP TO DAY OF HARVEST.**

### USES

**TANK MIXES:** SunSpray Ultra-Fine may be mixed with other pesticides to improve the level of kill or enhance coverage. Products with which SunSpray may be tank mixed include, but are not limited to, *Bacillus thuringiensis*, pyrethrum, malathion, and avermectin. Users should read and follow all directions and restrictions on the labels of the proposed tank mix products.

4 of 8

**LIMITED WARRANTY AND LIABILITY**

This product conforms to the description and is suitable for the uses set forth on this label but is strictly limited to those uses solely as directed under the specified label conditions, and only if purchased in the original unopened container. SELLER DISCLAIMS ANY AND ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THOSE RELATING TO MERCHANTABILITY AND FITNESS FOR PARTICULAR PURPOSE. Buyer and all users shall assume all risk,

liability and damage if this product is used, stored, handled or applied other than as specifically set forth on this label. SELLER'S LIABILITY AND BUYER'S OR USER'S REMEDIES SHALL BE LIMITED TO REFUND OF THE PURCHASE PRICE OR REPLACEMENT OF THE SPECIFIC QUANTITY INVOLVED. IN NO EVENT SHALL SELLER BE LIABLE FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

**TABLE 1A**

Crop	Pest	Application Rate - Gallons Oil			Time of Application (Stage of Development)	
		Dilute Spray per 100 Gal Water	Concentrate per 20-125 Gal. Water	Aerial* Min 20 Gal Spray		
Apples	Aphids (eggs)	2	6-8	6-8	Dormant or Delayed Dormant to 1/2" Green	
	Bugs (immature)	2	6-8	6-8		
	including Apple Red Bug					
	Mites (eggs) including European Red Mite	2	6-8	6-8		
	Scales (hard and soft)	2	6-8	6-8		
	Scurfy Scale	3	6-8	6-8		
	Fruit Tree Leaf Roller (eggs)	3	6-8	6-8		
	Mites	1	4	4	Summer (Foliar or Cover) or Postharvest Do not apply over fruit after waxy bloom forms	
Pears	European Red Mite (eggs)	1-2	4-6	4-6	Dormant or Delayed Dormant (up to and including petal fall)	
	Pear Leaf Blister Mite (eggs)	3	4-6	4-6		
	Pear Psylla	2	4-6	4-6		
	Scales	2	4-6	4-6		
	Fruit Tree Leaf Roller (eggs)	3	4-6	4-6		
		Two-Spotted Spider Mite	1	4	4-6	Summer (Foliar or Cover) or Postharvest
	Brown Mite	1	4	4-6		
	Pear Rust Mite	1-2	4-6	4-6		
	Pear Leaf Blister Mite	1.5-2	4-6	4-6		
	Pear Psylla	1.5-2	4-6	4-6		
Almonds Apricots Cherries	Aphids (eggs)	2-3	6-8	6-8	Dormant or Delayed Dormant	
	Bugs (immature)					
	Fruit Tree Leaf Roller (eggs)	2	4-6	4-6	Dormant	
	Tent Caterpillars					
San Jose Scale						
	Mites and Scales	1-1.5	4-6	4-6	Summer (Foliar or Cover) or Postharvest - Application should not be made over oil sensitive varieties - Do not apply to trees lacking adequate moisture	

\* Aerial application should be used only as emergency application when soil conditions do not permit regular ground application (helicopter only)

EPA Establishment No. 862-PA-1  
Net Contents: 2.5 Gallons

**BEST AVAILABLE COPY**

598

TABLE 1A (cont.)

Crop	Pest	Application Rate - Gallons Oil			Time of Application (Stage of Development)
		Dilute Spray per 100 Gal. Water	Concentrate per 20-125 Gal. Water	Aerial* Min. 20 Gal Spray	
Peaches Nectarine	Aphids (eggs) - Rugs (immature) Fruit Tree Leaf Roller (eggs) Mites (eggs) Peach Twig Borer (hibernaculae) Cottony Peach Scale San Jose Scale	3	6 - 8	6 - 8	Dormant or Delayed Dormant.
	Mites Scales	1 - 2	4 - 6	4 - 6	Summer (Foliar or Cover) or Postharvest.
Plums Prunes	Aphids (eggs) Scales Mites (eggs)	1.5 - 2	6 - 8	6 - 8	Dormant or Delayed Dormant to 1/2" Green Tip
	Mites Scales	1 - 1.5	4 - 6	4 - 6	Summer (Foliar or Cover) or Postharvest. Do not apply over certain fresh market fruits after bloom starts to form as the oil spray will remove the waxy bloom. Application should not be made over oil sensitive varieties.
Pecans	Aphids (eggs) Scales Mites (eggs)	3	6 - 8	6 - 8	Dormant
	Aphids Mites	1 - 1.5	4 - 6	4 - 6	Summer (Foliar or Cover) or Postharvest

\* Aerial application should be used only as emergency application when soil conditions do not permit regular ground application (helicopter only)

TABLE 1B

Crop	Pest	Application Rate - Gals Oil		Comments
		Dilute Spray per 100 Gal. Water	Low Volume* Spray per 80-125 Gals. Water	
Lemons and Grapefruit	Mites Scales including California Red Scale	1.0 - 1.4 1.0 - 1.6	10 - 20	For best results, spray during Fall, early Winter & Spring months when red scale are more easily controlled and trees generally less reactive to oil. For all citrus oil sprays ensure adequate soil moisture leaves should not wilt before noon
Navel* & Valencia Oranges, Other Citrus Varieties Including Non-Bearing	Mites including Rust Mite Spider Mite Scales including Black Scale California Red Scale Whitefly & Blackfly Sooty Mold	1.0 - 1.4 1.0 - 1.3 1.3 - 1.6 1.0 - 1.3 1.0 - 1.3	10 - 20	For low volume oil spray treatment during the Spring, application should be dis- continued as soon as tem- peratures go above 90° F during the day or relative humidity is expected to go down to 20 percent or below Oils applied after Oct. may increase cold damage to trees. Oils applied in Aug and Sept. may affect solids content and retard coloring The addition of a tank mix pesticide will improve perfor- mance against scales

\* Low volume applications require a minimum of 10 gallons of oil per acre for trees up to 10 feet in height plus 1 gallon per acre for each additional foot of average tree height. For navels, with the development of bloom, the amount of spray oil added to the 10 gallon-per-acre dosage for trees taller than 10 feet, in gallons per acre additional foot of average tree height, should be reduced to 3/4 gallon and then to 1/2 gallon at petal fall and postbloom through the June drop period.

**ACCEPTED**  
AUG 10 1994  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 862-23

**TABLE 2 Shade Trees, Shrubs, Ornamentals, Flower and Foliage Plants, Christmas Trees**

Crop or Plant	Pest	Application Rate Gallons Oil per 100 Gallon Water	Time of Application
Shade Trees* and shrubs including Conifers, Deciduous Broadleaf Evergreens & Woody Ornamentals	Aphids Adelgids Bugs (immature) Certain Caterpillars Eriophyid Mites Gall Mites Lace Bug Leaf Beetle Larvae Leafminer Mealybug Psyllids (immature) Sawfly (larvae) Scales (immature) Spider Mites Whiteflies (immature)	2 - 3	Winter Dormant Period as needed.
		1 - 3	Summer (Foliar or Cover) as needed.
Ornamental Trees*, shrubs along city streets, other rights-of-way including Conifers, Deciduous and Broadleaf Evergreens			
Flowering and Foliage plants including Roses and other Flowering shrubs		2 - 3	Winter Dormant Period as needed.
Foliage Ornamentals & Bedding Plants		1 - 2	Summer (Foliar or Cover) as needed.
Christmas Trees*	Aphids Adelgids Scale (soft & hard) (immature) Spider Mites	2 - 4	Winter
		1 - 3	Summer

*Oil removes the glaucous (blue) bloom from such evergreens as Colorado Blue Spruce and Winter Spruce. Always use lower dosage or test spray oil sensitive plants such as Cryptomeria, Smoke Tree, Chamaecyparis, Juniper, Japanese Holly and Spruce. Tendency toward sensitivity: Red Cedar and Douglas Fir. Do not spray Walnut foliage.*

**CAUTION:** Spray no more than 4 times during the growing season; Use a 2 week minimum application interval. Do not spray when buds have fully opened and shoot elongation is occurring. Do not spray when there is obvious moisture deficit in leaves, or the plant is under stress. For fall maintenance applications, reduce rate to 2 gallons oil per 100 gallons water and limit use only to: American Red Oak, Japanese Black Pine, Dogwood, Weeping Cherry, Cornelian Cherry, Hopsuckle, Norway Maple, Purple Leaf Plum. Keep away from open blooms-bleaching and spotting has been observed with the open blooms of certain plants.

**TABLE 3 Small Fruits**

Crop	Pest	Application Rate Gallons Oil per 100 Gallon Water	Time of Application
Blueberries	Mites	2	Foliage
	Scale (eggs)	2	Dormant
	Scales	3	
Strawberries	Aphids Mites	1	Dormant and Foliage

Crop	Pest	Application Rate - Gallons of Oil		Time of Application
		Dilute Spray	Concentrate Spray* (40-100 Gals. Spray per Acre)	
Grapes	Mealybug	4 - 6 Gal. with 200 to 300 Gal. water/acre	Not recommended	Dormant - A tank mix using an insecticide improves control.
	Mealybug	Not recommended	4	Spray when crawlers are active and exposed usually around early Summer (around July 1)
	Leathopper	Not recommended	4	Best results will occur when sprayed during first brood. Sprays after grapes are more than 1/4" diameter may affect bloom.

\* Air-carrier or air-blast type sprayers strongly recommended to avoid plant injury and reduce effect on bloom of table grapes.

**TABLE 4 Veg**

Crop
Asparagus
Beans
Beets
Cabbage
Cauliflower and other cole crops
Celery
Corn
Cucurbit
Eggplant
Lettuce
Melon
Pepper
Potato
Radish
Squash
Tomato
Tobacco

**TABLE 5 Field**

Crop
Corn (sweet & field)
Sugar Beets

**TABLE 4 Vegetables**

Crop	Pest	Application Rate Gallons Oil per 100 Gallon Water	Time of Application
Asparagus	Aphids	1-2	As Needed
Beans	Mites	1-2	
Beets	Beetle Larvae	1-2	
Cabbage	Leafminers	1-2	
Cauliflower and other cole crops	Certain Caterpillars	1-2	
Celery	Thrips	1-2	
Corn	Leafhopper	1-2	
Cucurbit	Whitefly	1-2	
Eggplant		1-2	
Lettuce		1-2	
Melon		1-2	
Pepper		1-2	
Potato		1-2	
Radish		1-2	
Squash		1-2	
Tomato		1-2	
Tobacco		1-2	

**TABLE 5 Field Crops**

Crop	Pest	Application Rate Gallons Oil per 100 Gallon Water	Time of Application
Corn (sweet & field) Sugar Beets	Aphids Mites Leafminers Certain Caterpillars including Corn Earworm, Rootworm and Armyworm Whitefly Bugs (immature)	2	As Needed

**TABLE 6 Greenhouse**

Crop or Plant	Pest	Application Rate Gallons Oil per 100 Gallon Water	Specific Comments
Azaleas	Aphids	1-2	Do Not Use on Coconut Palms or Maidenhair Ferns.
Begonias	Fungus Galls	1-2	
Camellias	Bugs	1-2	Chrysanthemum and Geranium blooms have shown Phytotoxic symp- toms at the 2 gallon rate
Chrysanthemum	Leafminers	1-2	
Crown of Thorns	Mealybugs	1-2	
Dieffenbachia*	Scales (soft & hard)	1-2	
Easter Lilies	Spider Mites	1	*Although no problems with Phytotoxicity have been seen at recommended rates, we recommend that the applicator conduct a Phytotoxicity test on 1 or 2 of the specific plants that are to be treated
Ferns	Whitefly (immature)	1-2	
Gardenias		1-2	
Geranium		1-2	
Hibiscus Foliage		1-2	
Jade Plant		1-2	
New Guinea Impatiens		1-2	
Palms		1-2	
Philodendron		1-2	
Poinsettias		1	
Portulaca		1-2	
Reiger Begonias		1-2	
Zinnias		1-2	
Leaf Pollah for hardy plants.		1	

*Frequency of Application: For the greenhouse pests listed, use once a week initially, then as the pest is controlled decrease the frequency to every 2 - 3 weeks as needed.*

*Application safety during bloom period should be determined for each individual species of plant to be treated by conducting a small test.*

**TABL**

Chrys  
Dieffe  
Draca  
Ferns  
Garde  
Jade P  
Palms  
Philoc  
Scheff

\* Protect l/c

Miscellar

Figs: Dor  
Mites, Me

Olives: P  
Apply at 4

Banana, f  
100 Gal. v  
venting it's

Avocado  
aphids, m

**BEST AVAILABLE COPY**

**LIMITED WARRANTY AND LIABILITY**

liability and damage if this product is used, other than as specified in the label.

Crop 8 9 8  
Page

**TABLE 7 Interiorscapes\***

Application Rate Gallons Oil per 100 Gallon Water	Specific Comments
1-2	Do Not Use on Coconut Palms or Maidenhair Ferns.
1-2	
1-2	Chrysanthemum and Geranium blooms have shown Phytotoxic symptoms at the 2 gallon rate.
1-2	
1-2	
1-2	
1	*Although no problems with Phytotoxicity have been seen at recommended rates, we recommend that the applicator conduct a Phytotoxicity test on 1 or 2 of the specific plants that are to be treated.
1-2	
1-2	
1-2	
1-2	
1-2	
1-2	
1-2	
1-2	
1-2	
1	

listed, use once a week initially, then as the pest is controlled.

needed for each individual species of plant to be treated by

Plant	Pest	Application Rate Gallons Oil per 100 Gallon Water	Comments
Chrysanthemum Dieffenbachia Dracaena Ferns Gardenias Jade Plant Palms Philodendron Schefflera	Aphids Leafminers Mealybugs Scales Spider Mites Whitefly (immature)	1-2	Do not apply to plants in direct sunlight behind glass. Do not use on Coccoloba Palms or Maidenhair Ferns. Chrysanthemum blooms have shown phytotoxic symptoms at the higher rate. Applicator should conduct a test for phytotoxicity by treating a few specimens before making a large scale application.

\* Protect floor, floor coverings and furnishings from overspray

**Miscellaneous:**

**Figs:** Dormant or Delayed Dormant: Fig Scale - Use 3 Gals. in 100 Gal. water as a foliar spray; Mites, Mealybug, Scale - Use 2 Gals. per 100 Gals. Water.

**Olives:** Postbloom through August and Postharvest: Scales - Use 1.5 gals. per 100 Gal. water. Apply at 400 to 800 Gal. per acre.

**Banana, Plantains:** Use as needed to control Yellow Sigatoka Disease - Use 1 - 1.5 Gal. per 100 Gal. water. This application is also effective in loosening Sooty Mold Fungus and in preventing it's formation by the control of Aphids, Mealybugs, Scales and Whitefly.

**Avocado (Haas Only) and Mangos:** Use 1 - 1.5 Gals. per 100 Gal. water as needed to control aphids, mealybugs, Scales and Whitefly.

	Aphids (eggs) Bugs (immature) including Apple Red Bug Mites (eggs) including European Red Mite Scales (hard and soft) Scurfy Scale Fruit Tree Leaf Roller (eggs)
	Mites
rs	European Red Mite (eggs) Pear Leaf Blister Mite (egg) Pear Psylla Scales Fruit Tree Leaf Roller (egg)
	Two-Spotted Spider Mite Brown Mite Pear Rust Mite Pear Leaf Blister Mite Pear Psylla
monds ricots berries	Aphids (eggs) Bugs (immature) Fruit Tree Leaf Roller Tent Caterpillars San Jose Scale
	Mites and Scales

\* Aerial application should be permit regular ground app

**BEST AVAILABLE COPY**