


865-15

PM-15

1/29/90

192



**HARRISON**

**VAPONA<sup>®</sup>\***  
*Kwik-Kil*  
**CATTLE  
SPRAY**

KEEP OUT OF THE REACH OF CHILDREN  
**WARNING!** SEE BACK PANEL FOR ADDITIONAL WARNINGS AND DIRECTIONS FOR USE  
HARRISON OIL CORP. MILWAUKEE, WISCONSIN  
NET CONTENTS 2.5 U.S. GALLONS

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.



# HARRISON

## 1% WAPONA® INSECTICIDE READY-TO-USE ANIMAL SPRAY SOLUTION

For the control of Horn Flies, Stable Flies, Face Flies, Houseflies and House Flies by direct application on dairy and beef cattle.

### KEEP OUT OF REACH OF CHILDREN WARNING

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. DO NOT GET IN EYES OR ON SKIN AND DO NOT BREATHE SPRAY MIST. Use safety goggles, respirator and body covering work clothes to avoid exposure. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. Keep away from open flames. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate food, water, or feedstuffs. If swallowed, do not induce vomiting and get medical attention immediately. Atropine and 2-PAM are antidotes.

**Physical and Chemical Hazards**  
Do not use or store near heat or open flame.

**Environmental Hazards**  
This product is toxic to fish, birds, and aquatic invertebrates. Do not apply directly to water or wetlands (swamps, marshes, bays, and saltmarsh). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Apply this product only as specified on this label.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in original container only. Protect from freezing. Do not store near heat or open flame.

**Container Disposal:** Triple-rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved State and local procedures.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray machine, or residue is a violation of Federal Law. If these wastes cannot be disposed of by use according to label directions, contact your State Pesticide or Environmental Control Agency, or the nearest Waste Management representative at the nearest EPA Regional Office for guidance.

**Warranty and Limitation of Damages**  
Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. Seller makes no other express or implied warranty, including any other express or implied warranty of fitness or of merchantability, and no agent of Seller is authorized to do so except in writing and with specific reference to this warranty. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material in which a claim is made.

WAPONA is a registered trademark of FENNER & SMITH ANIMAL HEALTH COMPANY.

### GENERAL DIRECTIONS

WAPONA® Horn-Kill Cattle Spray is a specially prepared ready-to-use insecticide for use on animals and hand, pump or automatic spray equipment, mist, fogging or misting equipment. Do not apply regularly to calves under 6 months of age. Sickened cattle should not be treated as they may show hypersensitivity to organic phosphorus. Do not treat animals within one day of slaughter. Do not apply directly in combination with Waterlox.

### USE AND DIRECTIONS FOR APPLICATION

**Direct Application to Dairy and Beef Cattle**  
(Horn Fly, Stable Fly, Face Fly, House Fly, Mosquitoes)

Apply only 1 to 2 full ounces of 1% solution per animal on a hot spot spray daily with hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply at intervals of 2 full ounces per day per animal. Use particular care to get over spray during extreme heat or when cattle are in a confined condition. No body-part should be necessary when used on beef animals. See precautionary warnings on this label.

**Spraying or Fogging of Animal Buildings**  
Horse Stables, Dairy Barns, Auction and Show Barns (Flies, Gnats, Mosquitoes)

Apply WAPONA® Horn-Kill Cattle Spray as a fog or mist. Remove all livestock from building before spraying. 1 pint of a 1% solution will treat 1000 cubic feet (45.23 m<sup>3</sup>). Before fogging reduce air movement as much as possible by closing doors, windows and other openings. Do not use this solution for fogging or misting pastures or areas where animals have received a direct application within 6 hours. Do not contaminate water, food or feedstuffs, milk, or milking utensils. See precautionary warnings on this label.

(EPA REG NO 895-15)  
(EPA EST NO 888-09-01)

FOR USE IN ALL TYPES OF FOGGERS AND SPRAYERS

### WARNING!

BEST AVAILABLE COPY