

DIRECTIONS

Fleas and lice — To control fleas and lice on dogs or cats, apply enough Green Light Flea and Tick Powder to treat animal thoroughly.

Ticks — Make heavier application for ticks, especially if engorged ticks are present.

If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary application, animals should be carefully examined for fleas, lice, or ticks before re-treating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

NOTE: Do not use on kittens under 4 weeks of age.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Do not breathe dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs, feed, feeding troughs, and watering receptacles.

KEEP OUT REACH OF CHILDREN

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

CONTAINER DISPOSAL: Crush and bury in non-crop areas away from streams or any body of water.

FLEA-TICK POWDER

ACCEPTED



Carbaryl (1-Naphthyl N-Methylcarbamate)* 95%
INERT INGREDIENTS 5%
TOTAL	100%

E.P.A. Reg. No. 869-5

Kills Fleas, Ticks, Lice!

GREEN LIGHT COMPANY

ACCEPTED

JUL 30 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 869-5 SUBJECT
TO ATTACHED COMMENTS.

869-5

GREEN LIGHT

FLEA *and* TICK POWDER

for **DOGS and CATS**

NET WEIGHT
4 OZ.



DIRECTIONS

Fleas and lice — To control fleas and lice on dogs or cats, apply enough Green Light Flea and Tick Powder to treat the animal thoroughly.

Ticks — Make heavier application for ticks, especially if engorged ticks are present.

If animal has free run, treatments may have to be made every second or third week. To avoid unnecessary application, animals should be carefully examined for fleas, lice, or ticks before re-treating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

NOTE: Do not use on kittens under 4 weeks of age.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Do not breathe dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs, feed, feeding troughs, and watering receptacles.

KEEP OUT REACH OF CHILDREN

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

CONTAINER DISPOSAL: Crush and bury in non-crop areas away from streams or any body of water.

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl-1-N-Methylcarbamate)*	5%
INERT INGREDIENTS	95%
TOTAL	100%

*Savin trademark Union Carbide Chem. Co.

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE ADDITIONAL CAUTIONS ON BACK.

E.P.A. Reg. No. 869-5

Kills Fleas, Ticks, Lice!

GREEN LIGHT COMPANY, San Antonio, Texas 78246

ACCEPTED

JUL 30 1974

UNITED STATES DEPARTMENT OF AGRICULTURE
 ANIMAL AND PLANT HEALTH INSPECTION SERVICE
 SAN ANTONIO, TEXAS

869-5