DIRECTIONS

For control of general insect feeders such as flea beetles, cucumber beetles, young mealy bugs, garden fleahoppers, aphids, thrips, beet webworms, and army worms feeding on Irish potatoes, beans, peas, cucumbers, beets, tomatoes, leaf and head Irtuce, spinach, and parsley.

For control of specific insect feeders such as:

Cabbage worms and young harlequin cabbage bugs (nymphs), feeding on cabbage, tender greens, cauliflower, broccoli, Swiss chard, Brussels sprouts, and kohlrabi.

Vegetable weevils feeding on carrots, turnips, beets, and radishes.

Mexican bean beetles on beans.

Bean leaf beetles and pea weevils on beans and peas.

Melon worms, young squash bugs (nymphs) and squash

vine borers feeding on squash, cucumbers, pumpkins,
and melons.

Leafhoppers feeding on potatoes and beans.

Potato beetles on potatoes, tomatoes and eggplants.

Small tomato fruitworms and small tomato hornworms on tomatoes.

Asparagus beetles on asparagus.

Tarnished plant bugs on tomatoes, beans, and peas.

Apply Green Light Vegetable Garden Dust lightly when insects first appear. Dust both tops or 3 bottoms of leaves. If properly applied, dust will be hardly visible on the plants. Repeat at weekly intervals until pests are controlled.

NOTE: Do not repeat application to asparagus within three days and do not apply within one day of harvest. Do not use within three days of harvest of broccoli, cabbage, head lettuce, Brussels sprouts, cauliflower, kohlrabi, radishes, beets, or turnips, and within one day of harvest of other vegetables except the vegetables listed below. Do not apply within 14 days of harvest of leaf lettuce, parsley, spinach, beet tops, turnip tops, Swiss chard.

CAUTION: Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash thoroughly after using. Keep out of reach of children.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equip-



FOR CONTROL OF MANY DESTRUCTIVE INSECT PESTS IN THE SMALL HOME GARDEN

Coursins Seviet and Retenent



Carbaryl (1-Naphthyl
N-Methylcarbamate)*
Rotenone

Other cube resins. 2%
INERT INGREDIENTS 90%

Total 100%

*Reg. trademark Union Carbide Chem. Co.
CAUTION: KEEP OUT OF REACH OF CHILDREN
FOR ADDITIONAL CAUTIONS SEE BACK PANEL

E.P.A. Reg. No. 869-15

ANTONIO A WARREN





For control of general insect feeders such as flea beetles, cucumber beetles, young mealy bugs, garden fleahoppers, aphids, thrips, beet webworms, and army worms feeding on Irish potatoes, beans, peas, cucumbers, beets, tomatoes, leaf and head lettuce, spinach, and pareley.

For control of specific in ect feeders such as:

Cabbage worms and young hartequin cabbage bugs (nymphs), feeding on cabbage, tender greens, cauliflower, broccoli, Swiss chard, Brussels sprouts, and

Vegetable weevils feeding on carrots, turnips, beets, and radishes.

Mexican bean beetles on beans.

Bean leaf beetles and pea weevils on beans and peas. Melon worms, young squash bugs (nymphs) and squash vine borers feeding on squash, cucumbers, pumpkins, and meloris.

Leafhoppers feeding on potatoes and beans.

Potato beetles on potatoes, tomatoes and eggplants.

Small tomato fruitworms and small tomato hornworms on tomatoes.

Asparagus beetles on asparagus.

Tarnished plant bugs on tomatoes, beans, and peas.

Apply Green Light Vegetable Garden Dust lightly when insects first appear. Dust both tops and bottoms of leaves. If properly applied, dust will be hardly visible on the plants. Repeat at weekly intervals until pests are controlled.

NOTE: Do not repeat application to asparagus within three days and do not apply within one day of harvest. Do not use within three days of harvest of broccoli, cabbage, head lettuce, Brussels sprouts, cauliflower, kohlrabi, radishes, beets, or turnips, and within one day of harvest of other vegetables except the vegetables listed below. Do not apply within 14 days of harvest of leaf lettuce. parsley, spinach, beet tops, turnip tops, Swiss chard.

CAUTION: Harmful it inhaled or sallswed Do not preside dust. Do not take intercally. Skin cortact may be that fill Ave disordard. We not berequely entire in the Keep to from bot children

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.



FOR CONTROL OF MANY DESTRUCTIVE INSECT PESTS IN THE SMALL HOME GARDEN

Contains Sevin* and Retenone

ACTIVE INGREDIENTS: Carbaryl (1-Naphthyl

N-Methylcarbamate)* Rotenone Other cube resins..... INERT INGREDIENTS

Total *Reg. trademark Union Carbide Chem. Co.

CAUTION: KEEP OUT OF REACH OF CHILDREN. FOR ADDITIONAL CAUTIONS SEE BACK PANEL

NET WEIGHT ONE POUND



