

JUL 15 1991

Mr. Jeff Lundke
Green Light Company
P.O. Box 17985
San Antonio, TX 78217

BEST AVAILABLE COPY

Dear Mr. Lundke:

Subject: Green Light Tomato & Vegetable Garden Dust
EPA Registration No. 869-15
Your Submission of a Draft Label With a Cover Letter Dated
September 8, 1989 in Response to the Registration Standard
for Rotenone

Enclosed is a stamped copy of the draft label you submitted. It will be acceptable with the following change. Revise the "Pesticide Disposal" statement to read "Leave unused pesticide in the original container. Securely wrap the original container in several layers of newspaper and discard in trash."

The Confidential Statement of Formula (CSF) dated June 25, 1986 will be kept in the file for this product. A revised CSF may be required if changes are made in labels for manufacturing-use products supplying the active ingredients for this product.

Sincerely yours,



Phil Wetton
Product Manager (18)
Insecticide-Rodenticide Branch
Registration Division (N7503C)

Enclosure

30360:1:Schroeder,C.Disk:KEYRIC:06/17/91:RW:ND:AM

CONCURRENCES

SEARCHED								
SERIALIZED								
INDEXED								
FILED								

OFFICIAL FILE COPY

GREEN LIGHT TOMATO & VEGETABLE GARDEN DUST

For Control Of Many Insect Pests In The Garden

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl methylcarbamate).....	7%
Rotenone.....	1%
Other Associated Resins.....	2%
INERT INGREDIENTS.....	90%
Total	100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

BEST AVAILABLE COPY

EPA Reg. No. 869-15

EPA Est. No. 869-TX-1

Formulated by Green Light Co., San Antonio, TX 78217

Net Wt. 1 lb.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED, call a physician immediately. IF INHALED, remove victim to fresh air. IF ON SKIN, wash with soap and water. IF IN EYES, rinse with plenty of water. Call a physician.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This pesticide is extremely toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, potholes). Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. PESTICIDE DISPOSAL: Unwanted pesticide material leave in original container. Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse container. Discard in trash.

ACCEPTED
with COMMENTS
in EPA Letter

JUN 25 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

Apply Green Light Tomato & Vegetable Garden Dust when insects first appear. Dust both top and bottom of leaves. Repeat at weekly intervals until pests are controlled.

FOR CONTROL OF THE FOLLOWING INSECTS:

Cabbage Worms and Harlequin bugs (nymphs) feeding on cabbage, tender greens, cauliflower, broccoli, swiss chard, brussels sprouts, and kohlrabi.

Vegetable Weevils feeding on carrots, turnips, beets, and radishes.

Mexican Bean Beetle on beans.

Bean Leaf Beetle and Pea Weevil on beans and peas.

Melon Worms, Squash Bugs (nymphs), and Squash Vine Borers feeding on squash, cucumbers, pumpkins, and melons.

Leafhoppers feeding on potatoes and beans.

Potato Beetles on potatoes, tomatoes, and eggplants.

Tomato Fruitworm and Tomato Hornworm on tomatoes.

Asparagus Beetles on asparagus.

NOTE: Do not repeat application to asparagus within one day of harvest. Do not use within three days of harvest of broccoli, cabbage, head lettuce, brussels sprouts, cauliflower, kohlrabi, radishes, beets or turnips, and within one day of harvest of other vegetables except the vegetables listed below. Do not apply within 14 days of harvest of leaf lettuce, spinach, beet tops, turnip tops, and swiss chard.