

GENERAL DIRECTIONS FOR USE
CAUTION: May cause irritation to skin, eyes, nose, and throat. Avoid contact with skin, eyes, and clothing. Avoid breathing vapors or dust. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating or smoking. Keep out of reach of children and domestic animals. Keep children and pets off treated areas until granules have been washed into the soil.

DO NOT CONTAMINATE FEED AND FOODSTUFFS.

ENVIRONMENTAL CAUTION: This product is toxic to wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

Apply granules when soil temperature at a depth of 6 to 8 inches is 40° to 80°F. Soil should be in good tilth and contain adequate moisture for good seed germination. Extremely wet or dry soils should not be treated. Application can be made before, during, or after planting according to directions only. Use only on crops listed on the label. Apply in accordance with directions for Nematode Killer Granules only. Use only on mineral soils. Granules exposed on soil surfaces will be hazardous to poultry. Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label's cautions and directions.

PREPLANTING APPLICATIONS: For best results, apply before each growing season of crop. For satisfactory nematode control, do not disturb the soil after application for 7 to 14 days on shallow rooted crops and 4 to 6 weeks on deep rooted crops.

Over-all: Apply granules evenly in rows 12 inches apart at a depth of 6 to 8 inches. Seal IMMEDIATELY with hoe and flood with water.

POSTPLANTING APPLICATION FOR ANNUAL CROPS: Apply granules as a band or side dressing at least 4 inches deep, as close to plants as possible without disturbing or damaging the roots. Seal treated area IMMEDIATELY and flood with water. For best results, apply during each growing season of crop.

POSTPLANTING APPLICATION FOR PERENNIAL OR ESTABLISHED PLANTS: Apply granules in 12-inch spacing 6 to 8 inches deep as close to plants as possible without injuring roots. Treat when necessary to prevent reinfestation. Be sure to treat areas where feeder roots are found. Seal treated area IMMEDIATELY and flood with water.

APPLICATION RATES: Tomatoes, Eggplants, Peppers (except bell peppers): Apply as only a preplanting treatment either at 2 ounces per 20 linear feet in the row or at 2 ounces per 20 square feet over-all.

Snapbeans, Okra, Dogwood: Apply as a pre-planting or post-planting treatment either at 2 ounces per 15 linear feet in the row or at 2 ounces per 15 square feet over-all.

Peach, Grape, Apricot, Fig, English Walnut, Periwinkle, Pansy, Daisy, Violet, Azalea, Boxwood, Rose, Gardenia, Holly: Apply as a pre-planting or post-planting treatment either at 4 ounces per 15 linear feet in the row or at 4 ounces per 15 square feet over-all.

NOTE: 2 heaping tablespoons full of Granules is 1 ounce.

Pour 2 or 4 ounces Granules in paper cup. Squeeze sides of cup and pour granules into trench of soil very, very lightly and apply at the rates mentioned above. Cover immediately. Pack soil down with hoe and flood with water.

LIMITATIONS: For Peaches, Grapes, Apricots, Figs, and English Walnuts, do not apply within 30 days of harvest. For Eggplant, Okra, Peppers, Snapbeans, and Tomatoes, do not feed plant remains to livestock. Do not space rows to be treated closer than 12 inches apart.

USE PRECAUTIONS: Do not apply this product to Tomatoes, Eggplants, and Peppers as a post-planting application. Do not store near seeds, food, or feedstuffs. Store leftover granules in original container with lid tight to prevent evaporation of pesticide.

Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.



ACCEPTED
1975
Under the Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 507-333

For use on certain vegetables
May be used to control plant
nematode damage to cert

ACTIVE INGREDIENT:
1,2-dibromo-3-chloro
Other halogenated C
INERT INGREDIENTS

CAUTION: KEEP
See

E.P.A. Est. No. 869-TX-1

Net

GREEN LIGHT COMPANY, San Antonio Texas 78217

GREEN LIGHT COMP

FOR USE
 thing. Avoid breathing vapors or dust. Harmful if swallowed. Wash thoroughly with soap
 animals. Keep children and pets off treated areas until granules have been washed into the

is killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur.

FOR USE
 good tillth and contain adequate moisture for good seed germination. Extremely wet or dry
 directions only. Use only on crops listed on the label. Apply in accordance with directions
 will be hazardous to poultry. Consult your State Agricultural Experiment Station or your
 and directions.

atisfactory nematode control, do not disturb the soil after application for 7 to 14 days on

LY with hoe and flood with water.
 at least 4 inches deep, as close to plants as possible without disturbing or damaging the
 wing season of crop.

12-inch spacing 6 to 8 inches deep as close to plants as possible without injuring roots.
 Seal treated area IMMEDIATELY and flood with water.

nting treatment either at 2 ounces per 20 linear feet in the row or at 2 ounces per 20 square

er 15 linear feet in the row or at 2 ounces per 15 square feet over-all.

se, Gardenia, Holly: Apply as a pre-planting or post-planting treatment either at 4 ounces

oil very, very lightly and apply at the rates mentioned above. Cover immediately. Pack soil

ays of harvest. For Eggplant, Okra, Peppers, Snapbeans, and Tomatoes, do not feed plant

anting application. Do not store near seeds, food, or feedstuffs. Store leftover granules in



GREEN LIGHT

NEMATODE KILLER

5% GRANULES

For use on certain vegetable, fruit, nut, flower, and shrub crops.
 May be used to control plant parasitic nematodes and to reduce
 nematode damage to certain plants.

ACTIVE INGREDIENTS:

1, 2-dibromo-3-chloropropane..... 5.00%
 Other halogenated C3 compounds..... .25%

INERT INGREDIENTS.....

94.75%
 Total 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN
 See additional cautions on back label

ACCEPTED

SEP 1 1976

Under the Federal Insecticide,
 Fungicide, and Rodenticide
 Act as amended, for the pesticide
 registered under EPA Reg. No. 869-TX-1

E.P.A. Est. No. 869-TX-1

Net Weight 1-1/2 lbs.

E.P.A. Reg. No. 869-63-AA

San Antonio Texas 78217

GREEN LIGHT COMPANY, San Antonio, Texas 78217