

DIRECTIONS FOR USE

ROSES: Black spot, powdery mildew, rust: 1½ tablespoons Fungicide per gallon water or 6 ounces per 16 gallons. Make first application when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout the season. Make thorough application to bottom and top of leaves.

CARNATIONS, SNAPDRAGONS, PHLOX, CHRYSANTHEMUMS, ZINNIAS, MARIGOLDS, ASTERS: Alternaria leaf spot, powdery mildew, anthracnose: 1½ tablespoons Fungicide per gallon water or 6 ounces per 16 gallons. Treat at least weekly during periods of rapid new growth and severe infestation and twice weekly during periods of rapid new growth and severe infestation caused by continuous rains or high humidity.

AZALEA cuttings: Damping off: 1½ tablespoons Fungicide per one gallon water. Allow to soak 15 to 30 minutes placing in rooting media.

CUCUMBERS, summer squash: Anthracnose, downy mildew, powdery mildew: 1 to 2 tablespoons per gallon water or 4 ounces in 8 to 16 gallons water and apply 200 gallons spray per acre. Make first application when first true leaves appear and repeat at weekly intervals until crop is harvested. Do not apply spray within 7 days of harvest.

GRAPES: Downy mildew, powdery mildew, black rot: Apply 1½ tablespoons Fungicide per gallon water or 6 ounces per 16 gallons water. Make first application just before bloom, repeat immediately after bloom, and repeat one to three applications at 7 to 10 day intervals. Apply up to 300 gallons per acre. Do not apply spray within 21 days of harvest.

Note: During extended periods of hot, dry weather, foliage injury to several varieties of grapes may occur.

CAUTION Avoid inhalation of dust or spray mist. Avoid prolonged or repeated contact with skin. Avoid contamination of feed or foodstuffs. Harmful if swallowed. Keep out of reach of children.

GREEN LIGHT COMPANY, SAN ANTONIO, TEXAS



CONTROLS: BLACK SPOT AND MILDEW

ALSO FOR FLOWERS AND CERTAIN VEGETABLES AND FRUIT DISEASES SUCH AS:
LEAF SPOT, RUST, ANTHRACNOSE,
AND CERTAIN OTHERS.

Active Ingredients:	
Folpet [N-(trichloromethylthio) pht. alimide]	42.00%
Dinitro (1-Methylheptyl) phenyl crotonate*	4.42%
Other nitrophenols and derivatives chiefly dinitro (1-Methylheptyl) phenol	0.40%
Inert ingredients	53.00%
Total	100.00%

*Karathane, trademark Rohm & Haas, U. S. Pat. No. 2,526,660

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTIONS ON BACK OF LABEL.**

U.S.D.A. Reg. No. 869.65

NET WEIGHT 6 OUNCES

DIRECTIONS FOR USE

ROSES: Black spot, powdery mildew, rust: 1 1/2 tablespoons Fungicide per gallon water or 6 ounces per 16 gallons. Make first application when young leaves first appear. Repeat at 7 to 10 day intervals or as necessary throughout the season. Make thorough application to bottom and top of leaves.

CARNATIONS, SNAPDRAGONS, PHLOX, CHRYSANTHEMUMS, ZINNIAS, MARIGOLDS, ASTERS: Alternaria leaf spot, powdery mildew, anthracnose: 1 1/2 tablespoons Fungicide per gallon water or 6 ounces per 16 gallons. Treat at least weekly during periods of rapid new growth and severe infestation and twice weekly during periods of rapid new growth and severe infestation caused by continuous rains or high humidity.

AZALEA cuttings: Damping off: 1 1/2 tablespoons Fungicide per one gallon water. Allow to soak 15 to 30 minutes placing in rooting media.

CUCUMBERS, summer squash: Anthracnose, downy mildew, powdery mildew: 1 to 2 tablespoons per gallon water or 4 ounces in 8 to 16 gallons water and apply 200 gallons spray per acre. Make first application when first true leaves appear and repeat at weekly intervals until crop is harvested. Do not apply spray within 7 days of harvest.

GRAPES: Downy mildew, powdery mildew, black rot: Apply 1 1/2 tablespoons Fungicide per gallon water or 6 ounces per 16 gallons water. Make first application just before bloom; repeat immediately after bloom and repeat one to three applications at 7 to 10 day intervals. Apply up to 300 gallons per acre. Do not apply spray within 7 days of harvest.

Note: During extended periods of hot, dry weather, sunburn injury to several varieties of grapes may occur.

Avoid inhalation. If dust or spray mist is inhaled, avoid prolonged or repeated contact with skin. Avoid contamination of food or feed stuffs. If swallowed, keep out of reach of children.

GREEN LIGHT COMPANY, SAN ANTONIO, TEXAS



ROSE FUNGICIDE

Contains Folpet and *Karathane
Sprayable Powder



Active Ingredients:	
Folpet (N-(trichloromethylthio) phthalimide)	42.00%
Dinitro (1-Methylheptyl) phenyl crotonate*	4.42%
Other nitrophenols and derivatives chiefly dinitro (1-Methylheptyl) phenol	0.49%
Inert Ingredients	53.09%
Total	100.00%

*Karathane, trademark Rohm & Haas, U. S. Pat. No. 2,526,660

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTIONS ON BACK OF LABEL.