

DIRECTIONS FOR USE

5] **LAWN PESTS:** For control of dog ticks, fleas, chinch bugs, earwigs, fall armyworms, leafhoppers, sod webworms, millipedes, use one level cupful Sevin 50% Powder in 15 to 20 gallons of water for each 500 square feet of established lawn area. For best results, mow lawn and apply immediately after rain or watering. When spraying for dog ticks or fleas, spray inside and outside of dog house and other areas where dogs sleep. Put fresh bedding in dog house after spraying. Repeat in 2 or 3 weeks if necessary. Spray dog at same time with 4 tablespoons Sevin per 1 gallon water.

6] **FLOWERS, SHRUBS, and SHADE TREES:** To control bagworm, box elder bug, leaf miner, blister beetle, elm leaf beetle, flea beetle, gypsy moth, Japanese beetle, June beetle, lace bug, leafhopper, leaf roller, mealy bug, mimosa webworm, plant bug, cicada, puss caterpillar, rose slug, scale insects, tent caterpillar, use two tablespoons Sevin per gallon water, or 1/2 pound Sevin in 25 gallons water, and spray plants thoroughly. Do not use on Boston ivy or Virginia creeper. Repeat weekly or as needed.

7] **VEGETABLE Insects:** Beans for bean beetles, stink bugs; cucumber and squash for melonworms, cucumber beetles, squash bugs; tomato, potato, eggplant and pepper for potato beetles, flea beetles, leafhoppers, stink bugs and grasshoppers; cabbage and broccoli for flea beetles, harlequin bug and cabbage caterpillars, use two tablespoons Sevin per one gallon water, or 8 ounces in 25 gallons water. Some leaf injury may occur on cucurbits if treatments are made when tender foliage is wet or in presence of prolonged high humidity. Repeat at 5 to 7 day intervals as needed. Do not apply spray to vegetables within 3 days of harvest.

8] **FRUIT Insects:** Peaches, apricots, plums for codling moth, oriental fruit moth, peach twig borer, plum curculio, San Jose scale, use 2 tablespoons Sevin in one gallon water, or 8 ounces in 25 gallons. Make first spray when one half petals have fallen. Repeat spray 3 or 4 times at 7 to 10 day intervals. Do not apply within 3 days of harvest.

9] **CITRUS Insects:** To control red scale, black scale, brown soft scale, and orange tortrix worms, use same dosage recommended for peaches. Apply for full coverage in normal spray schedules. Consult State Extension Specialists for proper timing of sprays. Do not apply within 5 days of harvest.

10] **GRAPES:** For leaf folder, leafhopper, leaf skeletonizer, grape berry moth, use 4 tablespoons Sevin per one gallon water. Make no application within one day of harvest. Consult State Extension Service Specialists for timing of sprays for grapes.

**CAUTION:** Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid breathing of spray. Skin contact may be harmful. Avoid skin contact. Wash hands after handling.

**Note for Physician:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food. Keep out of reach of children.

This product is highly toxic to bees exposed to direct treatment of residues on crops. Protection information may be obtained from your State Extension Service Specialists.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

Do not reuse container. Destroy when empty.

GREEN LIGHT  
**50% SEVIN\***

SPRAYABLE POWDER



*Kills*

**RESISTANT  
DOG TICKS and FLEAS**

WHEN USED AS DIRECTED

FOR THE CONTROL OF  
MANY INSECTS ON VEGETABLES, FRUITS  
ORNAMENTAL FLOWERS, SHRUBS,  
SHADE TREES, AND LAWNS

Active Ingredient :  
Carbaryl (1-Naphthyl N-Methylcarbamate) 50%  
Inert Ingredients ..... 50%

\*Reg. trademark Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate. U. S. Pat. No. 2,903,478

E.P.A. Reg. No. 869-66      NET WEIGHT 8 OUNCES

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

See additional cautions on back.

GREEN LIGHT COMPANY San Antonio, Texas 78246

GREEN LIGHT COMPANY San Antonio, Texas 78246

**LAWN PESTS:** For control of dog ticks, fleas, chinch bugs, earwigs, fall armyworms, leafhoppers, sod webworms, millipedes, use one level cupful Sevin 50% Powder in 15 to 20 gallons of water for each 500 square feet of established lawn area. For best results, mow lawn and apply immediately after rain or watering. When spraying for dog ticks or fleas, spray inside and outside of dog house and other areas where dogs sleep. Put fresh bedding in dog house after spraying. Repeat in 2 or 3 weeks if necessary. Spray dog at same time with 4 tablepoons Sevin per 1 gallon water.

**FLOWERS, SHRUBS, and SHADE TREES:** To control bagworm, box elder bug, leaf miner, blister beetle, elm leaf beetle, flea beetle, gypsy moth, Japanese beetle, June beetle, lace bug, leafhopper, leaf roller, mealy bug, mimosa webworm, plant bug, cicada, puss caterpillar, rose slug, scale insects, tent caterpillar, use two tablespoons Sevin per gallon water, or ½ pound Sevin in 25 gallons water, and spray plants thoroughly. Do not use on Boston ivy or Virginia creeper. Repeat weekly or as needed.

**VEGETABLE Insects:** Beans for bean beetles, stink bugs; cucumber and squash for melonworms, cucumber beetles, squash bugs; tomato, potato, eggplant and pepper for potato beetles, flea beetles, leafhoppers, stink bugs and grasshoppers; cabbage and broccoli for flea beetles, harlequin bug and cabbage caterpillars, use two tablespoons Sevin per one gallon water, or 8 ounces in 25 gallons water. Some leaf injury may occur on cucurbits if treatments are made when tender foliage is wet or in presence of prolonged high humidity. Repeat at 5 to 7 day intervals as needed. Do not apply spray to vegetables within 3 days of harvest.

**FRUIT Insects:** Peaches, apricots, plums for codling moth, oriental fruit moth, peach twig borer, plum curculio, San Jose scale, use 2 tablespoons Sevin in one gallon water, or 8 ounces in 25 gallons. Make first spray when one half petals have fallen. Repeat spray 3 or 4 times at 7 to 10 day intervals. Do not apply within 3 days of harvest.

**CITRUS Insects:** To control red scale, black scale, brown soft scale, and orange tortrix worms, use same dosage recommended for peaches. Apply for full coverage in normal spray schedules. Consult State Extension Specialists for proper timing of sprays. Do not apply within 5 days of harvest.

**GRAPES:** For leaf folder, leafhopper, leaf skeletonizer, grape berry moth, use 4 tablespoons Sevin per one gallon water. Make no application within one day of harvest. Consult State Extension Service Specialists for timing of sprays for grapes.

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid breathing of spray. Skin contact may be harmful. Avoid skin contact. Wash hands after handling.

**Note for Physician:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of food. Keep out of reach of children.

This product is highly toxic to bees exposed to direct treatment of residues on crops. Protection information may be obtained from your State Extension Service Specialists.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

Do not reuse container. Destroy when empty.

GREEN LIGHT

50% SEVIN\*

SPRAYABLE POWDER

RESISTANT  
DOG TICKS and FLEAS

WHEN USED AS DIRECTED

Active Ingredient :

Carbaryl (1-Naphthyl N-Methylcarbamate) 50%

Inert Ingredients ..... 50%

\*Reg. trademark Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate U.S. Pat. No. 2,903,478

E.P.A. Reg. No. 869-66

NET WEIGHT & OUNCES

CAUTION: KEEP OUT OF REACH OF CHILDREN.

See additional cautions on back.

GREEN LIGHT COMPANY San Antonio, Texas 78246

GREEN LIGHT COMPANY San Antonio, Texas 78246