THIS CARTON CONTAINS 10-8 OZ. BOXES (20 DOUBLE-DUTY ROSE FEEDINGS) ACCEPTED

GREEN LIGHT DOUBLE DUTY

## ROSE CARE

6-10-4 with SYSTEMIC

Feeds Roses and Kills Bugs Named Below with Same Treatment

PLANT E.P.A. EST. NO. 869-TX-1 E.P.A. REGISTRATION NO. 869-76



NET WT. 5 LB.

Supplies food for <u>roses</u>
for 4 to 6 weeks and protects the
roses from attack by aphids, thrips,
red spider, lace bugs, leafhoppers, and
white flies for 4 to 6 weeks.

SEE INSTRUCTIONS ON REVERSE SIDE FOR CARE OF ROSES

Easy to apply — just sprinkle on soil under rose bush and water down to roots.

AUG 23 154

- Easy measure for correct dosage.
- Will not affect beneficial insects.
- Insecticide cannot be washed away by watering or rain after it gets into plant.
- Acid forming for roses growing in neutral or alkaline soils.

GREEN LIGHT COMPAN' SAN ANTONIO, TEXAS 78246

THIS FIVE POUND CARTON TREATS UP TO 40 ROSE BUSHES

THIS CARTON CONTAINS 10-8 OZ. BOXES (20 DOUBLE-DUTY ROSE FEEDINGS)

### GREEN LIGHT

DOUBLE DUTY

# ROSE CARE

6-10-4 with SYSTEMIC

Feeds Roses and Kills Bugs Named Below with Same Treatment

PLANT E P.A. EST NO 869-TX-1

E.P.A. REGISTRATION

NO. 869-76



NET WT. 5 LB.

Supplies food for roses

for 4 to 6 weeks and protects the

roses from attack by aphids, thrips,

rad spiller, last lugs, trafforpers, and

white flies for 4 to 6 years

SEE DISTURBING ON PARTICION CARE OF BOSTS

Active Ingredient: U. S. Pat. No. 2,759,01	0
O, O-Diethyl S-[2-(ethylthio) ethyl] phosphorodithioate* 15	%
Inert Ingredients	
Guaranteed Analysis	
Total Nitrogen	%
Available Phosphoric Acid P2Os 10°	%
Soluble Potash K2O	
Total Primary Plant Foods 20°	%
Total Chlorine, not more than	
Primary Plant Foods derived from: Ammonium Sulfate, Diar	n-
monium Phosphate, and Muriate of Potash. Acid Forming	
WARNING: KEEP OUT OF REACH OF CHILDREN.	
SEE OTHER WARNING ON BACK AND LACH PACKAGE INSIDE.	

- Easy to apply just aprinkle on soil under rose bush and water down to roots.
- 3 Easy massure for correct desage.
- a Will not affect beneficial insects.
- Innesticials connect be mashed away by wedering arrain after it note mio plant.
- Acid forming for roses growing in neutral or alkaline rolls.

CONTROL SERVE CONTROL WAS

THIS FIVE POUND CALIFOR THEATS UP TO 40 HOSE DUSITES

#### TIPS ON ROSE CARE

#### For Strong, Vigorous Roses, and Large Beautiful Blooms

SOIL: This is truly the determining factor in successful rose growing. Roses should be planted in a mixture of 1/3 clay, 1/3 sand or sandy loam, and 1/3 organic matter (peat, leaf mold, compost, or well-rotted barnyard manure). This mixture should be twelve inches deep. If soil is predominately clay, the beds should be 6 to 12 inches above ground level for good drainage.

WHERE TO PLANT: Roses do better in full sun, or at least a half-day of sun and good air circulation, but not exposed to high winds.

WHEN TO PLANT: Plant bare-root roses in fall or spring. Plant canned roses any time of the year except winter.

#### HOW FAR APART TO PLANT:

6 to 8 feet for climbers

21/2 to 3 feet for bush roses

1 1/2 to 2 feet for polyantha or baby ramblers

PLANTING: In the south, plant roses so that the crown is just above ground level. In the north, plant so that crown is about 2 inches below ground level. Use Green Light Root Stimulator and Starter Solution to start roses growing.

PRUNING: After bare-rooted roses are planted, cut all canes back to 8 to 10 inches from ground level. In spring, as soon as all danger of frost is over, cut old rose canes back ten to 12 inches above ground level. Prune climbers in June only. Cut out all dead and weak canes on climbers, leaving 4 or 5 good canes.

MULCHING: Peat, cane pulp, pine needles, pine bark, or leaf mold four inches deep over the rose bed will keep out weeds, and grass, conserve moisture, protect roots from cold damage, and keep soil cooler in summer.

FEEDING ROSES: A special rose food such as Green Light Rose Care with Systemic helps produce vigorous, strong flowering roses and protects the roses from insects also. Follow directions on the package.

WATERING: The roots of roses must have a constant supply of water. Roses planted in the right kind of soil cannot be overwatered. The lack of water during summer months helps kill more roses than any other one thing. Unless there is an ample supply of rainfall, water the roses at least once a week and maybe twice a week during the hot, dry summer months. Soak the soil around roses 8 to 10 inches deep; and not the rose leaves.

DISEASE CONTROL: Use Green Light Rose Systemic as directed for control of blackspot on leaves; mildew on leaves, twigs and buds; and for rust disease on leaves.

#### **DIRECTIONS FOR USE**

Each box contains ½ lb. Rose Care with Systemic. There are 3 general classes of roses: bush roses, climbers, and small polyantha roses. Use ½ box Rose Care for bush roses, 1 box for climbers, and ¼ box for small polyantha roses.

Treat in spring when new foliage appears and repeat at 4-to-6 week intervals during growing season.

Water soil to a depth of 3 to 4 inches before applying Rose Care. Tear open at perforated tab and sprinkle amounts noted above on soil under bush. After Rose Care with Systemic is applied, water again slowly to percolate plant food down to root system.

Do not apply this Rose Care with Systemic insecticide to soil around feeding area of fruit or nut trees, vegetables or any food crop. Never feed roses just before frost. If Rose Care gets on leaves, wash it off to prevent burning.

WARNING: Poisoning may occur as a result of exposure. May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Do not get on skin, in eyes, or on clothing. Wear rubber gloves.

Do not contaminate feeds or foodstuffs. Keep out of reach of children. Wash thoroughly after handling. Keep children and pets off treated areas until product is worked into the soil and treated areas have dried completely.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label. Contains organic phosphate insecticide (inhibits cholines-

Antidote: atropine sulfate.

terase).

If swallowed induce vomiting. Call physician.

Conditions of use are beyond seller's control. Seller's liability is limited to product replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products use beyond claims on carton.

GREEN LIGHT COMPANY SAN ANTONIO, TEXAS 78246

Destroy 8-ounce box and outside container when empty.

SOIL: This is truly the determining factor in successful rose growing. Roses should be planted in a mixture of 1/3 clay, 1/3 sand or sandy loam, and 1/3 organic matter (peat, leaf mold, compost, or well-rotted barnyard manure). This mixture should be twelve inches deep. If soil is predominately clay, the beds should be 6 to 12 inches above ground level for good drainage.

WHERE TO PLANT: Roses do better in full sun, or at least a half-day of sun and good air circulation, but not exposed to high winds.

WHEN TO PLANT: Plant bare-root roses in fall or spring. Plant canned roses any time of the year except winter.

HOW FAR APART TO PLANT:

6 to 8 feet for climbers

21/2 to 3 feet for bush roses

1½ to 2 feet for polyantha or baby ramblers

PLANTING: In the south, plant roses so that the crown is just above ground level. In the north, plant so that crown is about 2 inches below ground level. Use Green Light Root Stimulator and Starter Solution to start roses growing.

PRUNING: After bare-rooted roses are planted, cut all canes back to 8 to 10 inches from ground level. In spring, as soon as all danger of frost is over, cut old rose canes back ten to 12 inches above ground level. Prune climbers in June only. Cut out all dead and weak canes on climbers, leaving 4 or 5 good canes.

MULCHIEL: Peat, cane pulp, pine needles, pine bark, or leaf mold four inches deep over the rose bed will keep out weeds, and grass, conserve moisture, protect roots from cold damage, and heep soil cooler in summer.

FEEDING ROSES: A special rose food such as Green Light Rose Care with Systemic helps produce vigorous, strong flowering roses and protects the roses from insects also. Follow directions on the package.

WATERING: The roots of roses must have a constant supply of water. Roses planted in the right kind of soil cannot be overwatered. The lack of water during summer months helps kill more roses than any other one thing. Unless there is an ample supply of rainfall, water the roses at least once a week and maybe twice a week during the hot, dry summer months. Soak the soil around roses 8 to 10 inches deep; and not the rose leaves.

DISEASE COMMON. Use Green Light Rose Systemic as directed for control of black-spot on leaves; mildew on leaves, twigs and buds; and for rust disease on leaves.

was in the state of the

Each box contains ½ lb. Rose Care with Systemic. There are 3 general classes of roces: bush roses, climbers, and small polyantha roses. Use ½ box Rose Care for bush roses, 1 box for climbers, and ¼ box for small polyantha roses.

Treat in spring when new foliage appears and repeat at 4-to-6 week intervals during growing season.

Water soil to a depth of 3 to 4 inches before applying Rose Care. Tear open at perforated tab and sprinkle amounts noted above on soil under bush. After Rose Care with Systemic is applied, water again slowly to percolate plant food down to root system.

Do not apply this Rose Care with Systemic insecticide to soil around feeding area of fruit or nut trees, vegetables or any food crop. Never feed roses just before frost. If Rose Care gets on leaves, wash it off to prevent burning.

WARNING: Poisoning may occur as a result of exposure. May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Do not get on skin, in eyes, or on clothing. Wear rubber gloves.

Do not contaminate feeds or foodstuffs. Keep out of reach of children. Wash thoroughly after handling. Keep children and pets off treated areas until roduct is worked into the soil and treated areas have died completely.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product and years specified on this label.

Contains organic phart . . . secticide (inhibits cholinesterase).

Antidote: atropine sulf.

If swallowed induce v ..., Call physician.

Conditions of use are E pand seller's control. Seller's liability is limited to product replacement or purchase price refund. Seller makes no warranty, express or implied, concerning products use beyond claims on carton.

GREEP THE IT COMPANY

Destroy 8 ounce box and outside container when empty