

DIRECTIONS FOR USE

USE THIS PRODUCT ONLY AS DIRECTED ON THIS LABEL.

GENERAL: The use of a regular protective spray program will minimize the risk of disease damage and can generally be accomplished with lower rates and less frequent fungicide applications. Once diseases have become established higher rates of fungicides and more frequent applications are required to bring them under control. This fungicide may be used with other Green Light insecticides.

Lawn & Turf Grasses: As a protective spray or before disease is well established such as Brown Patch (*Rhizoctonia*), Melting Out (*Helmintosporium*), Red Thread (pink Patch), Copper Spot, Slime Mold (black sooty mold), and Rust on stem, crown and leaf, apply 4 ounces in 15 gallons of water per 1000 square feet at 14-day intervals.

When conditions are favorable for diseases use 6 to 8 ounces Fungicide in 15 gallons of water per 1000 sq. ft. at 7-day intervals.

For Pythium Blight, Dollar Spot (*Sclerotinia*), as a protective spray, use 6 to 8 ounces of Fungicide in 15 gallons of water at 6-day intervals. Continue until disease threat is past.

For Snow Mold (*Fusarium*) use 6 to 8 ounces Fungicide in 15 gallons of water and apply during winter at 2 to 6-week intervals.

Water lawn or turf thoroughly before treating with Fungicide and then do not water for several days thereafter.

NOTE: Do not graze treated areas. Do not feed clippings to livestock.

Ornamental Diseases: Mix 1-1/2 tablespoons Fungicide in 1 gallon of water and make full coverage spray. Spray when plants are well leafed out and apply at 7 to 10-day intervals throughout the season.

Black Spot on Roses: Use above mixture. Begin when plants are well leafed out and apply at weekly intervals during growing season.

Leaf and Flower Spot on Gladiolus caused by *Curvularia* and *Botrytis*: Before flower spikes appear in full coverage, spray at weekly intervals before disease appears and increase to 2 or 3 applications per week after diseases appear. Do not apply after spikes appear.

Leaf Spot on Iris, Carnation, Hollyhock: Start when plants are well leafed out and repeat at weekly intervals. Also for *Botrytis* blight on dahlia, lily, and tulips.

For Anthracnose and Rust on hollyhock, pansies, flowering dogwood and for Scab, Cedar Apple Rust on ornamental crabapples.

Petal Blight on Azalia, camellia, rhododendron, spray 2 to 3 times a week while flowers are opening. Direct spray into flowers and spray ground under bushes thoroughly.

Fruit Trees: Scab, Cedar-apple Rust, Black Rot, Fly Speck, Sooty Blotch, Bitter Rot, Brown Rot: Apples, Pears, Crabapples, Quince: Use 1 pound Fungicide in 50 gallons of water. Make application every 7 to 14 days throughout the season starting with green tip or delayed dormant spray and continue through cover sprays. Do not spray within 30 days of harvest. Do not graze livestock in treated areas. Consult local or State Agriculture Extension Agent for spray schedules and timing and number of sprayings.

Black Rot, Downy Mildew—Grapes: Use 1-1/2 pounds Fungicide in sufficient water (50 to 60 gallons) to provide thorough coverage. Start spraying when new shoots are 1/2 to 1-1/2 inches long. Repeat when shoots are 3 to 5 inches long and when shoots are 8 to 10 inches long, and continue at 14 day intervals, but do not apply Fungicide within 66 days of harvest.

NOTE: 2 tablespoons per gallon of water is equivalent to 1 pound of Fungicide per 50 gallons of water.

Vegetables: Tomatoes: Late and Early Blight, Gray Leaf Spots, Gray Leaf Mold, Anthracnose—Use 1 pound Fungicide per 50 gallons of water and spray plants sufficiently to provide thorough coverage. Start spraying when seedlings emerge and repeat at 7-day intervals throughout the season. Do not spray within 5 days of harvest.

Potatoes: Early and Late Blight—Use same dilutions as for tomatoes. Begin spraying when plants are 4 to 6 inches high and repeat at 7 to 10 days and throughout the season.

Cucumbers, Summer Squash, Cantaloupe, Watermelons: Downy Mildew, Anthracnose, Scab, Ceraspora Leaf Spot, Gummy Stem Blight—Use same dilutions as for tomatoes and spray thoroughly. Start spraying when plants are in two-leaf stage and repeat as for tomatoes. Make sure both upper and lower leaf surfaces are covered with spray. Do not spray within 5 days of harvest.

ENVIRONMENTAL CAUTION: This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes.

IMPORTANT: Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness and may also generate flammable vapors. Keep container closed when not in use. Destroy when empty by burning or crushing and burying in an unoccupied area away from lakes or streams. If burned, stay out of smoke.

KEEP OUT OF REACH OF CHILDREN.



ACCEPTED
SEP 27 1976
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 869-84

Controls the Spot, Slime Mold Blight, Anthracnose, Speck, Sooty Blotch, Leaf Spot.

See directions on Active Ingredient A co-ordinator in which the in Mangar Zinc + Ethylene

INERT INGREDIENTS

MAY CAUSE IRRITATION OF SKIN CONTACT ATTENTION. IF SWALLOWED

E.P.A. Est. No. 8

Distributed by **GREEN LIGHT COMPANY**, San Antonio, Texas 78217

Distributed by

ON THIS LABEL.
 Damage and can generally accomplish with lower rates and less frequent applications are required to bring them
 down Patch (Rhizoctonia), Melting Out (Helminthosporium), Red
 and leaf, apply 4 ounces in 15 gallons of water per 1000 square
 ft. per 1000 sq. ft. at 7-day intervals.
 Fungicide in 15 gallons of water at 6-day intervals. Continue until
 ing winter at 2 to 6-week intervals.
 14 days thereafter.

coverage spray. Spray when plants are well leafed out and apply
 weekly intervals during growing season.
 weekly intervals. Also for Botrytis blight on dahlia, lily, and tulips.
 le Rust on ornamental crabapples.
 opening. Direct spray into flowers and spray ground under bushes

Rot: Apples, Pears, Crabapples, Quince: Use 1 pound Fungicide
 ing with green tip or delayed dormant spray and continue through
 ees. Consult local or State Agriculture Extension Agent for spray

60 gallons) to provide thorough coverage. Start spraying when
 d when shoots are 8 to 10 inches long, and continue at 7 to 14
 as of water.

Use—Use 1 pound Fungicide per 50 gallons of water and spray
 id repeat at 7-day intervals throughout the season. Do not spray
 when plants are 4 to 6 inches high and repeat at 7 to 10 days and

e, Scab, Ceraspore Leaf Spot, Gummy Stem Blight—Use same
 itage and repeat as for tomatoes. Make sure both upper and lower

or ponds. Do not contaminate water by cleaning equipment or
 become wet or overheated in storage. This may bring on chemical
 ions. Keep container closed when not in use.
 lakes or streams. If burned, stay out of smoke.

ACCEPTED
SEP 27 1976
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under
 EPA Reg. No. 869-84

GREEN LIGHT
GENERAL PURPOSE
FUNGICIDE
FOR CONTROL OF MANY DISEASES ON LAWN GRASSES, FRUITS,
VEGETABLES AND ORNAMENTAL PLANTS.

**CONTROLS THESE PLANT DISEASES: Downy Mildew, Melting Out, Copper
 Spot, Scab, Rust, Anthracnose, Leaf Spot, Black Spot, Petal Blight, Black Rot, Fly
 Speck, Sooty Blotch, Litter Rot, Brown Rot, Downy Mildew, Early & Late Blight, Alternaria
 Leaf Spot.**

See directions on back for Lawn Grasses, Ornamentals, Fruits, Vegetables.

Active Ingredients:
 A co-ordination product of Zinc ion and Manganese Ethylene Bisdithiocarbamate 80%
 in which the ingredients are

Manganese++	16%
Zinc++	2%
Ethylene bisdithiocarbamate ion (C ₄ H ₆ N ₂ S ₄)	62%
INERT INGREDIENTS	20%
Total	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
MAY CAUSE IRRITATION OF NOSE, THROAT, EYES, AND SKIN. AVOID BREATHING SPRAY MIST. IN CASE
OF SKIN CONTACT WASH WITH WATER, FOR EYES WASH COPIOUSLY WITH WATER AND GET MEDICAL
ATTENTION. IF SWALLOWED GIVE TABLESPOON SALT IN GLASS OF WARM WATER. CALL PHYSICIAN.
SEE OTHER PRECAUTIONS ON BACK PANEL

E.P.A. Est. No. 869-TX-1 NET WEIGHT 1 POUND E.P.A. Reg. No. 869-84-ZA

San Antonio, Texas 78217

Distributed by GREEN LIGHT COMPANY, San Antonio, Texas 78217