ACTIVE INGREDIENTS:

Beta-naphthoxyacetic acid 0.004% iNERT INGREDIENTS 99.996%

Total 100.00%

During period of cool nights when temperature gets below 55° F., the tomato blooms are not pollenized and thus will not set fruit.

DIRECTIONS FOR USE

Shake well before using.

Remove cap from aerosol can. Hold can 18 to 20 inches from tomato blooms. Spray backs of flowers in cluster. Avoid getting excessive amount of spray on tomato foliage. Only a small amount of spray need be put on the blooms. About one second of spraying per tomato bloom cluster is sufficient. Apply to each flower cluster once when most of flowers in cluster are open but not more than three applications per plant per season.

NOTE: Do not apply to tomatoes within 15 days of harvest.

CAUTION: Avoid excessive dosage of spray to tomato flowers or leaves. Do not use on other vegetables or fruits. Do not breathe spray mist. May be harmful if swallowed. Wash hands with soap and water after using and before smoking. Keep out of reach of children.





NET U

NET CONTENTS 13 AV. OZ. U.S.D.A. Reg. No. 869-93

GREEN LIGHT COMPANY San Antonio, Texas

AN AID TO:
INCREASING TOMATO SET
INCREASING SIZE OF TOMATO

SPRAY WHEN 3 OR 4 TOMATO FLOWERS IN CLUSTER ARE OPEN. REPEAT SPRAY AT WEEKLY INTERVALS DURING PERIOD OF COOL NIGHTS.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTIONS ON BACK