

**PRE-HARVEST USE LIMITATION:**

NO LIMITATION on Beans, Blueberries, Carrots, Cucumbers, Eggplants, Melons, Grapes, Peppers, Pumpkins, Squash, Tomatoes, and Okra.

ALLOW 1 DAY between dusting and harvest of Cranberries and Strawberries.

ALLOW 3 DAYS between dusting and harvest of Cabbage, Cauliflower, Brussels Sprouts, and Broccoli.

**DIRECTIONS: READ ENTIRE LABEL BEFORE USING.**

Use sufficient dust to obtain full coverage, being sure to cover upper and lower leaf surfaces. Apply when insects or their damage appear. Repeat at 7 to 10-day intervals or as necessary. Some plants and the insects controlled on them are:

**GRAPES:** Grape leaf folder, grape leafhopper and grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetle, and red-banded leaf roller.

**STRAWBERRIES:** For control of meadow spittlebug, strawberry leaf roller, and strawberry weevil.

**BLUEBERRIES:** Blueberry maggot, cherry and cranberry fruitworms, and Japanese beetle.

**CRANBERRIES:** Cutworms, fireworms, fruitworms, Japanese beetle, and leafhoppers.

**BEANS:** (Green, Lima, snap beans, cowpeas, and blackeyed peas) — Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhopper, velvetbean caterpillar, and western bean cutworm, armyworms, stink bugs, and tarnish plant bug.

**CUCUMBER, MELONS, PUMPKIN AND SQUASH:** Pickleworm, melonworm, cucumber beetle, flea beetle, leafhopper, and squash bugs. May cause some leaf injury during prolonged high humidity. Do not use on watermelons in Florida.

**TOMATO, EGGPLANT, AND PEPPER:** Colorado potato beetle, flea beetle, and leafhopper, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnish plant bugs, and stink bugs.

**OKRA:** Stink bugs. Repeat at 5 to 7-day intervals.

**CARROTS:** Flea beetles, harlequin bug and leafhopper, six-spotted leafhopper (aster yellows vector), armyworm, spittlebug, tarnish plant bug, and stink bugs.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER:** Flea beetle, harlequin bug, and armyworm.

**DOGS AND CATS:** Ticks and Fleas. Dust liberally over entire animal, avoiding eyes; rub thoroughly into skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Also, dust liberally in and around sleeping quarters, kennels, and doghouses. Be sure to force dust into cracks and crevices. Repeat as necessary. Avoid treatment of kittens and puppies less than 4 weeks old.

**LAWN INSECTS:** Dust liberally using 1 lb. per 275 sq. ft. for control of ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworm (lawn moths), millipedes. For best results mow lawn and apply after watering or rain. Repeat in 2 to 3 weeks if necessary.

**NOTE:** Does not control spider mites but is compatible with all common miticides.

**CAUTION:** Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs, and watering receptacles.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**CONTAINER DISPOSAL:** Crush and bury in non-crop areas away from streams or any body of water.

**GREEN LIGHT**



**5% SEVIN\* DUST**

**FOR CONTROL OF INSECT PESTS**

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate)\* ..... 5%

INERT INGREDIENTS ..... 95%

\*Sevin trademark of Union Carbide Corp.  
U.S. Patent No. 2,903,478

**CAUTION: KEEP OUT OF REACH OF CHILDREN.  
SEE OTHER CAUTIONS ON BACK PANEL.**

**NET WT. 1 POUND**

U.S.D.A. Reg. No. 869-118

Formulated by **GREEN LIGHT COMPANY** San Antonio, Texas 78246

869-118

11-22-1971

1/2

EPH  
11-10-71

ACCEPTED  
NOV 22 1971  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO.....SUBJECT  
TO ATTACHED COMMENTS.

869-118