

Read entire label before using this Dust.

DIRECTIONS FOR USE

VEGETABLES: Cover upper and lower surfaces of the leaves thoroughly.

Potato, Tomato, Eggplant, Pepper: To control Colorado potato and flea beetles, leafhoppers, tomato fruitworm, fall armyworm, tomato hornworm, stink bugs, lace bugs, tarnished plant bugs. May be used on day of harvest.

Okra: To control corn earworm, stink bugs. May be used on day of harvest.

Cucumbers, Squash, Pumpkin, Melons: To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, squash bugs. May be used on day of harvest.

Leafy Vegetables and Root Crops such as Carrots, Chinese Cabbage, Collards, Dandelion, Endive, Beet, Horseradish, Kale, Lettuce, Mustard greens, Parsley, Parsnip, Radish, Rutabaga, Salsify, Spinach, Swiss chard, Turnip: To control flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, cabbage loopers, imported cabbage worms, tarnished plant bugs, stink bugs. May be used on day of harvest of carrots, up to 3 days before harvest of other root crops and head lettuce, and up to 14 days before harvest of other leaf crops.

FLOWERS: To control Japanese, blister, flea, and June beetles; boxelder, mealy, lace, and plant bugs; leafhoppers; leaf rollers; psyllid; rose aphid; and exposed thrips.

NOTE: Does not control spider mites but is compatible with all common miticides.

SMALL FRUITS: Cover upper and lower surfaces of the leaves thoroughly.

Cane Berries such as Blackberry, Boysenberry, Dewberry, Loganberry, Raspberry, to control Japanese beetle, leaf rollers, leafhoppers, raspberry aphid. Can be used up to 7 days before harvest.

Strawberries for spittlebugs, leafhoppers, and weevils, to be used up to 1 day before harvest.

LAWNS: Dust liberally using one pound per 275 sq. ft. for control of ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworm (lawn moths), millipedes. For best results mow lawn and apply after watering or rain. Repeat in 2 to 3 weeks if necessary.

DOGS AND CATS: Dust liberally over entire animal, avoiding eyes, to control ticks and fleas. Rub thoroughly into skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Also, dust liberally in and around sleeping quarters, kennels, and doghouses. Be sure to force dust into cracks and crevices. Repeat as necessary. Avoid treatment of kittens and puppies less than 4 weeks old.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs, feed, feeding troughs, and watering receptacles.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of ponds, lakes, and streams.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

CONTAINER DISPOSAL: Crush and bury in non-crop areas away from streams or any body of water.

GREEN LIGHT



5% SEVIN^{*} DUST

ACCEPTED
APR 16 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 869-118 - SUBJECT
TO ATTACHED COMMENTS

For Vegetables, Flowers, Lawns, Small Fruits,
Also Dogs and Cats To Control Many Insects
Feeding On These Plants and Pets

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate)* 5%
INERT INGREDIENTS 95%
*Sevin trademark of Union Carbide Corp.
U.S. Patent No. 2,903,478

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER CAUTIONS ON BACK PANEL.

NET WEIGHT 1 POUND

E.P.A. Reg. No. 869-118

Formulated by **GREEN LIGHT COMPANY** San Antonio, Texas 78246

869-118

04/16/1973

1/1