

PM 9

869-118

1 of 2

GREEN LIGHT SEVIN® 5% DUST

TO CONTROL MANY INSECTS FEEDING ON VEGETABLES, FLOWERS, FRUITS, LAWNS, DOGS AND CATS.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl methylcarbamate).....	5%
INERT INGREDIENTS.....	95%
Total	100%

ACCEPTED
JUL 14 1995
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 869-118

Sevin® registered trademark of Rhone-Poulenc for Carbaryl Insecticide

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 869-118

EPA Est. No. 869-TX-1

Formulated by Green Light Co., P.O. Box 17985, San Antonio, TX 78217-0985

Net Weight 1 lb. (454 gm)

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing vapors (dust of spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately. IF INHALED, remove victim to fresh air and apply respiration if indicated. IF ON SKIN, wash promptly with soap and water. Rinse thoroughly. IF IN EYES, rinse eyes for at least 15 minutes with water and call a physician immediately.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment of residues on blooming crops or weeds. if bees are visiting the treatment area. **DO NOT APPLY TO CARPETS AS STAINING MAY OCCUR.**

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **DO NOT USE INDOORS.**

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. **PESTICIDE DISPOSAL:** Leave unwanted pesticide material in original container. **CONTAINER DISPOSAL:** Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

129

VEGETABLES: Cover upper and lower surface of the leaves thoroughly. **POTATO, TOMATO, EGGPLANT, PEPPER:** To control Colorado potato and flea beetles, leafhoppers, tomato fruitworm, fall armyworm, tomato hornworm, stink bugs, lace bugs, tarnished plant bugs. May be used on day of harvest. **OKRA:** To control corn earworm, stink bugs. May be used on day of harvest. **CUCUMBER, SQUASH, PUMPKIN, MELONS:** To control pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, squash bugs. May be used on day of harvest. **LEAFY VEGETABLES and ROOT CROPS** such as **CARROTS, CHINESE CABBAGE, COLLARDS, DANDELION, ENDIVE, BEET, HORSERADISH, KALE, LETTUCE, MUSTARD GREENS, PARSLEY, PARSNIP, RADISH, RUTABAGAS, SATSIFY, SPINACH, SWISS CHARD, TURNIP:** To control flea beetles, leafhoppers, harlequin bug, six-spotted leafhopper (aster yellows vector), corn earworm, armyworms, cabbage loopers, imported cabbageworms, tarnished plant bugs, stink bugs. May be used on day of harvest of carrots, up to 3 days before harvest of other root crops and head lettuce, and up to 14 days before harvest of other leaf crops.

FLOWERS: To control Japanese, blister, flea, and June beetles, boxelder, mealy, lace and plant bugs, leafhoppers, leafrollers, psyllid, rose aphid and exposed thrips. **NOTE:** Does not control spider mites, but is compatible with all common miticides.

SMALL FRUITS: Cover upper and lower surfaces of leaves thoroughly. Cane berries such as: **BLACKBERRY, BOYSENBERRY, DEWBERRY, LOGANBERRY, RASPBERRY** to control Japanese beetle, leafrollers, leafhoppers, raspberry aphid. Can be used up to 7 days before harvest. **STRAWBERRIES** for spittlebugs, leafhoppers and weevils. To be used up to 1 day before harvest.

LAWNS: Dust using 1 lb. per 275 sq.ft. for control of ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworm (lawn moth), millipedes. For best results mow lawn and apply after watering or rain. Repeat in 2 to 3 weeks if necessary.

DOGS AND CATS: Dust entire animal, avoiding eyes, to control fleas and ticks. Rub thoroughly into skin. Repeat at weekly intervals if needed. Also, dust in and around sleeping quarters, kennels, and doghouses. Be sure to force dust into cracks and crevices. Repeat as necessary. Avoid treatment of puppies and kittens less than 4 weeks old. Do not use on pregnant pets. Do not allow recently treated pets on carpet as staining may occur.