04 PM UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

DEC | 5 | 1999

Jeff Luedke Green Light Company P.O. Box 17985 San Antonio, Texas 78217-0985

Subject:

Green Light Sevin 5% Dust

EPA Registration No. 869-118

Amendment to delete unsupported uses and revise label as

per GDE instructions dated August 6, 1999.

Amendment dated October 22, 1999.

Dear Mr. Luedke:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, Rodenticide Act, as amended, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

1. Correct the PHIs for Cucumber, Squash, Pumpkin, Melons, and Okra from 7 days to 3 days. Correct the PHIs for Mustard greens from 7 days to 14 days. Correct the PHIs for Chinese Cabbage, Collards, and Kale from 3 days to 14 days.

Submit two copies of your final printed labeling before you release the product for shipment.

Sincerely,

George Tompkins, Ph.D., Entomologist Insecticide-Rodenticide Branch Registration Division (7505C)

READ ENTIRE LABEL BEFORE EACH USE.
USE ONLY ON DOGS OR CATS.
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing vapors (dust spray or mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

FIRST AID: IF SWALLOWED: Drink one or two glasses of water and Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately. IF INHALED: Remove victim to fresh air and apply respiration if indicated. IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. IF IN EYES: Rinse eyes for at least 15 minutes with water and call a physician immediately.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- . Change clothing as soon as possible after use.
- Wash the outside of gloves before removing.
 As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consuit a veterinarian immediately.

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. However, flea and tick collars may be immediately replaced.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds if bees are visiting in the treatment area.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. DO NOT USE INDOORS. FOR HOME USE ONLY.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. PESTICIDE DISPOSAL: Unwanted pesticide material leave in original container in several layers of newspapers and discard in trash. CONTAINER DISPOSAL: Do not reuse container. Discard in trash.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a lig-

uid carbaryl prode on not repeat application more than once a week.

VEGETABLES: Cover upper and lower surface of the leaves thoroughly. Potato, Tomato, Eggplant, Pepper: To control Colorado Potato and Flea beetles, Leafhoppers, Lace bugs, Tomato fruit worm, Fall armyworm, Tomato hornworm, Stink bugs, Lace bugs, Tarnished plant bugs. May be used up to 7 days of harvest of Potato; up to 3 days of harvest of Tomato, Eggplant and Pepper. Okra: To control Corn earworm, Stink bugs. May be used up to 7 days of harvest. Cucumber, Squash, Pumpkin, Melons: To control Pickleworm, Melonworm, Cucumber beetles. Leafhoppers, Squash bugs. May be used up to 7 days of harvest.

Leafy Vegetables and Root Crops such as Carrots, Chinese Cabbage, Collards, Dandellon, Endlve, Beet, Horseradish, Kale, Lettuce, Mustard greens, Parsley, Parsnip, Radish, Rutabaga, Salsify, Spinach, Swiss chard, Turnip: To control Flea beetles, Leafhoppers, Harlequin bug, Six-spotted leafhopper (aster yellows vector), Corn earworm, Armyworms, Cabbage loopers, Imported Cabbage worms, Tarnished plant bugs, Stink bugs, May be used up to 7 days of harvest of carrots, horseradish, radish, parsnip, rutabaga, salsify, mustard greens, beets (roots), turnips (roots); up to 3 days before harvest of chinese cabbage, collards, kale, and up to 14 days before harvest of lettuce, endive, dandelion, parsley, spinach, swiss chard, beets (tops), turnips (tops).

FLOWERS: To control Japanese, Blister, Flea, and June beetles; Boxelder, Mealy, Lace, and Plant bugs; Leathoppers, Leaf rollers, Psyllid; Rose aphid; and Exposed thrips.

Note: Does not control Spider mites but is compatible with all common miticides.

SMALL FRUITS: Cover upper and lower surfaces of leaves thoroughly. Cane Berries such as Blackberry, Boysenberry, Dewberry, Loganberry, Raspberry, to control Japanese beetle, Leaf rollers, Leafhoppers, Raspberry aphid. Can be used up to 7 days before harvest.

Strawberries for Spittlebugs, Leafhoppers, and Weevils, to be used up to 7 day before harvest. LAWNS: Dust liberally using one pound per 275 sq. ft, for control of Ants, Chinch bugs, Earwigs, Fall armyworm, Fleas, Leafhoppers, Mosquitoes, Sod webworm (Lawn moths), Millipedes. For best results mow lawn and apply after watering or rain, Repeat in 2 to 3 weeks if necessary.

DOGS AND CATS: To control Ticks and Fleas. Use product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Reapply every week. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puoples under 12 weeks of age. Dust bedding and doghouse using a shaker can or other dust applicator. Also, dust in and around sleeping quarters, kennels, and doghouses. Be sure to force dust into cracks and crevices. Repeat as necessary but not more than once a week. Do not allow recently treated pets on carpet as staining may occur.

EPA Est, No. 869-TX-1 EPA P

EPA Reg. No. 869-118

COPYRIGHT © GREEN LIGHT CO.
GREEN LIGHT COMPANY, P. O. BOX 17985
SAN ANTONIO, TX 78217-0985



To Control Many Insects Feeding On Vegetables, Flowers, Lawns, Fruits, Dogs and Cats.

ACTIVE INGREDIENT:

*Sevin registered trademark of Rhône-Poulenc Total 100% for Carbaryl Insecticide

KEEP OUT OF REACH OF CHILDREN
See Back Panel For Additional
Precautionary Statements





Net Weight 1 lb. (454 gm)