

READ ENTIRE LABEL BEFORE USING

To control sowbugs (pill bugs), crickets, snails, darkling ground beetles, cutworms, armyworms, earwigs, and grasshoppers in the lawn, flower bed, ornamental, and home vegetable gardens, follow these instructions carefully.

DIRECTIONS FOR USE

Punch holes in top of canister and sprinkle bait LIGHTLY on the ground where pests are found, giving particular attention to edges of walks, flower beds, vegetable gardens, and along foundation of house and porch. Best results will be obtained if the bait is applied in late afternoon or evening because most of the above pest feed late in the evening and at night. For best results, water area to be treated before applying bait. Do not put bait out in piles. Scatter it so the bait can hardly be seen on the ground or grass.

This bait is stomach poison and works slowly. Results will be apparent in 1 to 3 days after applying. Repeat the application at four or five-day intervals until satisfactory control is obtained. One pound of bait will treat approximately 500 square feet except for vegetable gardens.

VEGETABLE GARDENS: On beans, eggplant, potatoes, tomatoes, cabbage, lettuce, beets, turnips, spinach, to control cutworms, armyworms, crickets, sowbugs, snails, use at rate of 40 pounds per acre or for home gardens apply on ground between rows one pound of bait per 1100 square feet or 11 rows 100 feet long with a band treatment of bait 1 foot wide. Avoid getting bait on vegetables. Do not use bait within 3 days of harvest around cabbage, lettuce, beets, turnips, nor within 14 days of harvest around beet tops, leaf lettuce, spinach and turnip tops.

WARNING May be fatal if swallowed, inhaled, or absorbed through skin. Avoid breathing harmful dust or spray mist. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking. Wear clean clothing. Avoid contamination of feed and foodstuffs.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL CAUTIONS: This product is toxic to birds and other wildlife. Birds feeding on treated areas may be killed. Treated bait exposed on soil surface may be hazardous to birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSAL OF CONTAINER: Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.



**GREEN LIGHT
Ready-to-use Bait**

**BUG BAIT
AND
SNAIL BAIT**

**FOR USE IN LAWNS,
GREENHOUSES,
ORNAMENTAL GARDENS,
HOME VEGETABLE GARDENS**

Controls

**THIS IS A MEAL BAIT,
NOT PELLETS**

**One pound of bait
treats approximately 500 sq. ft.**

• **SOW BUGS**
(Pill bugs)

• **EARWIGS**

• **CRICKETS**

• **SNAILS**

• **GRASSHOPPERS**

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl methycarbamate)*	5%
Metaldchyde	1%
INERT INGREDIENTS	94%
Total	100%
*Sevin trademark of Union Carbide Corp. U.S. Pat. No. 2,903,478 and 3,009,855	

E.P.A. Est. No. 869-TX-1 NET WEIGHT - one pound

**WARNING: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER WARNINGS ON PACK.**

**WARNING: THIS PESTICIDE MAY BE FATAL TO CHILDREN, DOGS
OR OTHER PETS IF EATEN. KEEP CHILDREN AND PETS OUT OF
THE TREATED AREA. FOLLOW DIRECTIONS AND USE WITH CARE.**

READ ENTIRE LABEL BEFORE USING

To control sowbugs (pill bugs), crickets, snails, darkling ground beetles, cutworms, armyworms, earwigs, and grasshoppers in the lawn, flower bed, ornamental, and home vegetable gardens, follow these instructions carefully.

DIRECTIONS FOR USE

Punch holes in top of canister and sprinkle bait LIGHTLY on the ground where pests are found, giving particular attention to edges of walks, flower beds, vegetable gardens, and along foundation of house and porch. Best results will be obtained if the bait is applied in late afternoon or evening because most of the above pests feed late in the evening and at night. For best results, water area to be treated before applying bait. Do not put bait out in piles. Scatter it so the bait can hardly be seen on the ground or grass.

This bait is stomach poison and works slowly. Results will be apparent in 1 to 3 days after applying. Repeat the application at four or five-day intervals until satisfactory control is obtained. One pound of bait will treat approximately 500 square feet except for vegetable gardens.

VEGETABLE GARDENS: On beans, eggplant, potatoes, tomatoes, cabbage, lettuce, beets, turnips, spinach, to control cutworms, armyworms, crickets, sowbugs, snails, use at rate of 40 pounds per acre or for home gardens apply on ground between rows one pound of bait per 1100 square feet or 11 rows 100 feet long with a band treatment of bait 1 foot wide. Avoid getting bait on vegetables. Do not use bait within 3 days of harvest around cabbage, lettuce, beets, turnips, nor within 14 days of harvest around beet tops, leaf lettuce, spinach and turnip tops.

WARNING: May be fatal if swallowed, inhaled, or absorbed through skin. Avoid breathing harmful dust or spray mist. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking. Wear clean clothing. Avoid contamination of feed and foodstuffs.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL CAUTIONS: This product is toxic to birds and other wildlife. Birds feeding on treated areas may be killed. Treated bait exposed on soil surface may be hazardous to birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSAL OF CONTAINER: Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.



**GREEN LIGHT
Ready-to-use Bait**

**BUG BAIT
AND
SNAIL BAIT**

Controls

**FOR USE IN LAWNS,
GREENHOUSES,
ORNAMENTAL GARDENS,
HOME VEGETABLE GARDENS**

**THIS IS A MEAL BAIT,
NOT PELLETS**

• **SOW BUGS**
(Pill bugs)

One pound of bait
treats approximately 500 sq. ft.



• **EARWIGS**




• **CRICKETS**



• **SNAILS**



• **GRASSHOPPERS**



E.P.A. Est. No. 869-TX-1

NET WEIGHT - one pound

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl methylcarbamate)*	5%
Metaldhyde	1%
INERT INGREDIENTS	94%
Total	100%
*Sevin trademark of Union Carbide Corp. U.S. Pat. No. 2,903,478 and 3,009,855	

**WARNING: KEEP OUT OF REACH OF CHILDREN.
SEE OTHER WARNINGS ON BACK.**

**WARNING: THIS PESTICIDE MAY BE FATAL TO CHILDREN, DOGS
OR OTHER PETS IF EATEN. KEEP CHILDREN AND PETS OUT OF
THE TREATED AREA. FOLLOW DIRECTIONS AND USE WITH CARE.**