

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

Jeff Luedke Green Light Company P.O. Box 17985 San Antonio, Texas 78217-0985 OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

APR 4 2000

Subject:

Green Light Bug Bait and Snail Bait

EPA Registration No. 869-119

Amendment dated January 12, 2000 in response to EPA

Letter dated December 23, 1999 in response to GDE.

Dear Mr. Luedke:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended labeling:

1. Change the application rate for the crop beans so that the application rate does not exceed one pound of bait per 1500 square feet or 15 rows 100 feet long. The maximum application rate specified in the carbaryl GDE for beans is 1.5 lbs ai/acre per application.

Submit two copies of your final printed labeling before you release the product for shipment. A copy of the labeling stamped "Accepted With Comments" is enclosed for your records.

Sincerely,

George Tompkins, Ph.D., Entomologist Insecticide-Rodenticide Branch Registration Division (7505C)

ACCEPTED with COMMENTS In EPA Letter Dated:

APR 4 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 269-119

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking. Wear clean clothing. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately. IF INHALED: Remove person from dust area. Call a loian. IF ON SKIN: Wash promptly with soap and water.

IF IN EYES: Flush eyes with cold water. Get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treatment area. DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. To control sowbugs, pillbugs, crickets, snails, slugs, cutworms, armyworms, earwigs, and grasshoppers in the lawn, flower beds, ornamental and home vegetable garden, sprinkle bait LIGHTLY on the ground where pests are found, giving particular attention to edges of walks, flower beds, vegetable gardens, and along foundation of house and porch. Best results will be obtained if the bait is applied in late afternoon or evening because most of the above pests feed late in the evening and at night. For best results, water area to be treated before applying bait. Do not put bait out in piles. Scatter it so the bait can hardly be seen on the ground or grass. The bait is stomach poi-'d works slowly. Results will be apparent in 1 to 3 days after applying. Ithe application at four or five-day intervals until satisfactory control is obtained. One pound of bait will treat approximately 500 square feet except for vegetable gardens.

VEGETABLE GARDENS: On beans, eggplant, potatoes, tomatoes, cabbage, lettuce, beets, turnips, spinach and strawberries, to control cutworms, armyworms, crickets, sowbugs, snails. For home gardens apply on ground between rows 1 pound of bait per 1100 square feet or 11 rows 100 feet long with a band treatment of bait 1 foot wide. Avoid getting bait on vegetables. Do not use bait within 3 days of harvest around cabbage, beans, tomatoes, eggplant; within 7 days of harvest around potatoes; nor within 14 days of harvest around lettuce, leaf lettuce, beets, beet tops, turnips, turnip tops, spinach and strawberries.

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Store in original container. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

EPA Reg. No. 869-119 EPA Est. No. 869-TX-1

FORMULATED BY GREEN LIGHT COMPANY P.O. BOX 17985, SAN ANTONIO, TX 78217-0985







For Use in Lawns, Flower Beds, Ornamental and Home Vegetable Gardens.

This is a Meal Bait, not Pellets. One pound of bait treats approximately 500 sq ft.

approximately 500 sq ft.

GRASSHOPPERS/CRICKETS


ACTIVE INGREDIENTS:

tetraoxycyclo-octane) ...1%
INERT INGREDIENTS: ...94%

TOTAL 100%



SOW BUGS/ PILL BUGS



CUTWORMS

KEEP OUT OF REACH OF CHILDREN CAUTION See Back Panel For Additional Precautionary Statements.

NET WT. 1 LB. (454gm)