5-13-1982

869-123

WEEDS CONTROLLED

Green Light Broadleaf Weed Killer is especially effective against weeds such as Sheep Sorrel, Knotweed, Spotted Spurge, Chickweed, Oxalis, and Henbit. In addition to these, it kills virtually all broadleaf weeds, some of which are listed below.

Bedstraw	Cocklebur	Lambsquarter	Poison Ivy	Spurge
Black Medic	Dandelion	Lespedeza	Poison Oak	Velvet Leaf
Buckhorn	Dock	Mallow	Purslane	Wild Carrot
Bull Thistle	Galinsoga	Morning Glory	Ragweed	Wild Garlic
Burdock	Ground Ivy	Mustard	Sheep Sorrel	Wild Lettuce
Buttercup	Hawkweed	Oxalis	Shepherd's Purse	Wild Onion
Chicory	Heal-all	Peppergrass	Speedwell	Yarrow
Chickweed	Henbit	Pigweed	Tansy Mustard	Many other broadleaf
Clover	Knotweed	Plantains	Thistle	weeds

DIRECTIONS FOR USE

GENERAL INFORMATION: Broadleaf Weed Killer is specially formulated for protection to seeding grasses and 2, 4-D sensitive grasses such as St. Augustine grass, bentgrass, and buffalo grass. Sprays should be applied when broadleaf weeds are small and tender for best results. If weeds are older and hard to kill the higher rates recommended should be used. Avoid spraying St. Augustine grass under conditions of high temperature and moisture when the St. Augustine is growing rapidly. Under these conditions, or in cases of accidental overdose, some slight temporary discoloration of the St. Augustine foliage might occur. DO NOT USE ON DICHONDRA LAWNS. Do not use on newly seeded lawns. Do not contaminate water intended for irrigation or domestic purposes.

To ensure weeds are growing and able to absorb the herbicide, irrigate the lawn 2 or 3 days prior to spraying. Do not water for 24 hours after treatment. Although spray works faster with temperatures above 70° F., sprays can be applied at any time plants are growing. Spray when air is calm to avoid drift. Spray will injure or kill clover, dichondra, and most ornamental and vegetable crops. This product works slowly. Therefore it will be from 1 to 2 weeks before weeds die. Amount of water in spray is not important except to evenly distribute the chemical. If directions call for one ounce in 1 gallon on 300 square feet, it may be more convenient to use one ounce in 2 gallons on 300 square feet, depending on the sprayer. Generally, large volumes of water result in more uniform coverage.

HOW TO MIX:

1. For use on St. Augustine grass, bamgrass, buffalo grass and grass seedlings mix 1 ounce or 2 tablespoons in 1 gallon of water and uniformly spray 300 square feet. (Example: an area 10 feet wide and 30 feet long). If necessary, repeat treatment in 2 or 3 weeks, if hard-to-kill weeds persist.

2. For use on all other grasses such as bermuda, bahia, blue grass, centipede, fescue, rye, mix 1-1/2 ounces in 1 gallon of water and spray uniformly over 300 square feet. On hard to kill species, or under poor growing conditions, or on older weeds, mix 2 ounces in 1 gallon of water and spray on 300 square feet with special attention to thorough coverage of hard-to-kill weeds.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. If contact with eyes occurs, flush with plenty of water.

Prevent drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not contaminate domestic or irrigation water. Do not store in contact with fertilizers, seeds, insecticides, or fungicides. Use only lawn sprayers for this use. Coarse sprays are less likely to drift. Do not apply to areas underlaid by roots of desirable trees and shrubs.

When container is empty, wash thoroughly and destroy. Never re-use. Destroy by perforating or crushing and burying in a safe place. Do not burn.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label when conditions of use—such as weather, compatibility with other chemicals, and condition of application equipment—will vary, Seller makes no claims other than those stated on this label.

Distributed by GREEN LIGHT COMPANY, San Antonio, Texas 78217

GREEN LIGHT

WEED & FEED

KEEP FROM FREEZING

CONTAINS 2, 4-D, MCPP, AND DICAMBA

ONE PINT COVERS UP TO 4,800 SQUARE FEET

ACTIVE INGREDIENTS:	
Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid*	7.23%
Dimethylamine salt of 2, 4-Dichlorophenoxyacetic acid**	2.07%
Dimethylamine salt of Dicamba (3, 6-dichloro-o-anisic acid)***	.88%
INERT INGREDIENTS	89.82%
Contains Trimec (R)†	Total 100.00%

*2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.60 lbs. per gallon.

**2, 4-Dichlorophenoxyacetic acid equivalent 0.14 lbs. per gallon.

***3, 6-dichloro-o-anisic acid equivalent 0.08 lbs. per gallon.

†Registered trademark of PBI Gordon Corporation U.S. Pat. No. 3,284,186

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on back

Distributed by GREEN LIGHT COMPANY, San Antonio, Texas 78217

ACCEPTED

MAY 13 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 869-123