

N-89-76

869-125

CAUTION:

Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after using. Keep away from fire or sparks. In case of contact, flush skin or eyes with plenty of water, -for eyes, get medical attention.

IMPORTANT: Never allow Green Light Systemic Fungicide to become wet during storage. This may lead to certain chemical changes which will reduce its effectiveness as a fungicide. Keep container closed when not in use. Do not reuse container. Bury when empty.

DIRECTIONS FOR USE

Green Light Systemic Fungicide is recommended for the control of certain diseases of roses. It is highly effective on roses, providing both curative (eradicator) action and protective action. Green Light Systemic Fungicide is a wettable powder to be mixed with water for application as a spray.

ROSES (FIELD AND GREENHOUSE) . . . Powdery Mildew, Black Spot: Use one tablespoonful per two (2) gallons of water. Begin application when disease first appears and repeat at 10 to 14 day intervals throughout the growing season; shorten intervals during humid, rainy weather.

HOME FRUITS: Green Light Systemic Fungicide is recommended on peaches, nectarines, apricots, plums, to control brown rot, peach scab.

BROWN ROT: Dilute 2 ounces of Fungicide in 12½ gallons of water. Spray trees (in combination with dormant oil) as a delayed dormant treatment prior to bud break. Use 1 ounce in 12½ gallons of water at early bloom (popcorn, red bud, or green tip) and at full bloom.

Beginning 3 weeks before harvest, apply 1 to 2 additional sprays of 1 ounce Fungicide per 12½ gallons of water.

PEACH SCAB: Use same schedule as for brown rot plus applications of 1 ounce Fungicide in 12½ gallons of water at shuck-split and shuck-fall.

Spray trees thoroughly to the point of runoff. Do not graze livestock in treated orchards.

HOME GARDEN VEGETABLES: Tomatoes for white and gray mold, leaf spot, use 1 ounce Systemic Fungicide in 12½ gallons water.

For cucumbers, squash for powdery mildew, anthracnose, use ½ ounce Systemic Fungicide per 12½ gallon water.

Begin application when diseases first appear. Repeat spray at 7 to 14 day intervals as needed. Spray plants thoroughly.

LAWN DISEASES: For control of brown patch, dollar spot, use 1 ounce per 1000 sq. ft. on a preventive schedule and 2 ounces per 1000 sq. ft. at a curative rate with sufficient water to obtain uniform coverage (2-5 gallons per 1000 sq. ft.). As a preventive begin at the earliest appearance of disease and continue at 10-day to 14-day intervals throughout the season. At the curative rate apply at the first appearance of disease and repeat as needed. When conditions are unusually favorable for development of large brown patch and the disease is active, reduce the interval between sprays to 5-7 days. Do not graze treated areas or feed clippings to livestock.

Addition of a spreader-sticker to the spray mixture enhances curative action of the fungicide, and improves distribution of the spray on hard-to-wet plants such as roses.

Do not reuse container. Bury when empty.