

PM 21 869-125

10/3

GREEN LIGHT SYSTEMIC FUNGICIDE
Disease Control with Benomyl

For Control of Leafspots, Mildews, Blights, Molds, Scab and Other Diseases on Vegetables, Fruits, Nuts, and Lawns.

ACTIVE INGREDIENT:

Benomyl [Methyl 1-(butylcarbamoyl)-2-benzimidazole carbamate]	50%
INERT INGREDIENTS	50%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 869-125

EPA Est. No. 869-TX-1

Distributed by Green Light Co., San Antonio, TX 78212-0985

Net Wt. 2 oz.

ACCEPTED
JUN 13 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 869-125

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May irritate eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and skin. Wash thoroughly after using. **FIRST AID:** In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention. This product may cause a temporary allergic reaction in a few susceptible persons. This condition should be treated as an allergic dermatitis. There is no evidence of aftereffects or permanent injury. If sensitivity occurs, protective clothing should be worn during periods of exposure such as during application or when contacting treated foliage. **PROTECTIVE MEASURES:** Always spray with your back to the wind. Wear long-sleeved shirt, long pants and rubber gloves. Wash gloves thoroughly with soap and water before removing. Do not enter or allow reentry into treated areas until sprays have dried unless wearing the protective clothing specified under "Protective Measures".

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when runoff is likely to occur. Drift and runoff from treated areas may be hazardous to fish in adjacent areas. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated.

PHYSICAL OR CHEMICAL HAZARDS: Keep away from fire or sparks.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Never allow to become wet during storage. This may lead to certain chemical changes which will reduce its effectiveness. Keep container closed when not in use. Keep in original container. **DISPOSAL:** Securely wrap original container. Rinse thoroughly before discarding in trash.

Green Light Systemic Fungicide is recommended for control of many important plant diseases. It is highly effective, providing both curative (eradicant) and protective action. Green Light Systemic Fungicide is ready to be mixed with water for application as a spray. Spray plant surfaces sufficiently to provide thorough coverage. Use higher application rates where indicated when disease is active and conditions favor further development.

VEGETABLES: Tomatoes - For control of Gray Mold (Botrytis), Leaf Mold (Cladosporium), White Mold (Sclerotinia), Cercospora Leaf Spot, Phoma Leaf Spot - Apply 1/2 to 1 Tablespoon of Systemic Fungicide per 1 gallon of water. Begin applications when disease first appears and repeat at 7 to 14 day intervals as needed. May be used up to day of harvest.

Cucumbers, Melons, and Squash - For control of Target Spot (Cucumbers), Gummy Stem Blight, Powdery Mildew, Anthracnose - Apply 1/2 Tablespoon per gallon of water. Begin applications when plants begin to run or when diseases first appears, and repeat at 7 to 14 day intervals as needed. May be used up to day of harvest.

Beans - For control of White Mold (Sclerotinia), Gray Mold (Botrytis) - Apply 1 to 2 Tablespoons per gallon. Repeat applications 7 to 10 days apart as needed. Do not apply within 14 days of harvest.

FRUITS: Strawberries - For control of Gray Mold (Botrytis), Powdery Mildew, Leaf Scorch, Leaf Blight, Mycosphaerella Leafspot - Apply 1 Tablespoon per gallon of water. Repeat applications at 10 to 14 day intervals.

NUTS: Pecans - For control of Pecan Scab, Brown Leafspot, Downy Spot, Powdery Mildew, Liverspot, Zonate Leafspot, Fungal Leaf Scorch - Apply 1/2 to 1 Tablespoon per gallon. Apply at prepollination when young leaves are unfolding, when small nuts are forming, and thereafter at 3 to 4 week intervals. Do not apply after shuck split.

LAWNS: For control of Brown Patch, Dollar Spot, use 2 ounces per 1000 sq. ft. on a preventive or curative schedule with sufficient water to obtain uniform coverage (2 to 5 gallons per 1000 sq. ft.). As a preventive, begin at the earliest appearance of

30/3

disease and continue at 10 to 14 day intervals throughout the season. As a curative, apply at the first appearance of disease and repeat as needed. When conditions are unusually favorable for development of Large Brown Patch and the disease is active, reduce the interval between sprays to 5 to 7 days. Do not graze treated areas; do not feed clippings to livestock.

CC
CC
CC

CC
CC
CC
CC
CC
CC
CC
CC
CC
CC