

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)**

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin. In case of contact with skin, wash with soap and water. Avoid breathing spray mist. Do not contaminate feed or drinking water. **NOTE FOR PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidote.


ENVIRONMENTAL HAZARDS: This product is highly toxic to bees exposed to direct treatment or residues on crops. This product is toxic to fish. **KEEP** out of any body of water. Apply this product only as specified on this label. Protective information may be obtained from your Cooperative Agricultural Extension Service. **DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.**

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Keep in original container. Do not store near heat or open flame.

PESTICIDE DISPOSAL: Unwanted pesticide material, leave in original container. Securely wrap original container in several layers of news paper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.



STATEMENT OF PRACTICAL TREATMENT IF SWALLOWED: Drink one of two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything to eat or drink unless a physician is present. **IF INHALED:** Remove victim from area and apply respiration if indicated. **ON SKIN:** Wash promptly with soap and water. Rinse thoroughly. **IF IN EYES:** May irritate eyes, nose, throat and skin. Rinse eyes for at least 15 minutes with water and call a physician immediately.

Read entire label before using. Use only as directed and follow cautions carefully.

SHAKE WELL BEFORE POURING OUT OF BOTTLE

DILUTION: Use 4 teaspoons per 1 gallon water or 1-1/3 pints per 32 gallons water, except on lawns. Spray all plants thoroughly covering underside and tops of leaves and branches.

SHADE TREES, SHRUBS, FLOWERS: Japanese beetle, flea and June beetles, boxelder weevil, lace, and plant bugs, leaf hoppers, leafrollers, psyllid, rose aphid and spotted thrips, spider mite, bagworms, scale insects, mimosa webworm, tent and puss caterpillars, cicada, orange tortra.

VEGETABLE INSECTS: Tomato, potato, eggplant and pepper for Colorado potato beetle, flea beetles, lace bugs, tomato hornworm, Beans for Mexican bean beetle, bean leaf beetle, cucumber beetles, Japanese beetle, leafhoppers, cucumber, melons, pumpkin and squash for pickleworm, melonworm, cucumber beetle, squash bugs. Apply 1 gallon spray per 535 sq. ft. Spray may be used within 1 day of harvest.

FRUIT INSECTS: Apples and pears for codling moth, green apple aphid, Japanese beetles, leafroller, bagworm, periodical cicada, San Jose scale, crape killers, pear psylla.

NOTE: Sevin Spray will cause thinning of fruit on many apple varieties when applied 10 to 25 days after full bloom. Consult your local County Agricultural Extension Agent for varieties affected.

PEACHES: Plums, apricots, nectarines for codling moth, plum curculionid, oriental fruit moth, Japanese beetle, leafroller, periodical cicada. Apply 1 gallon spray per 535 sq. ft. Spray may be used within 3 days of harvest. Apply spray at petals fall and every 7 to 14 days thereafter until controls are checked.

NOTE: Proper timing of the sprays for control of fruit insects is essential. Consult your local Agricultural Extension Service for proper fruit tree spray schedule.

LAWN INSECTS: Ground bugs, fire ants, armyworms, leafhoppers and weevils, and other grass pests. Mix 2 quarts use 10 ounces per 20 gallons of water per 100 square feet or 1 ounce per 2 gallons of water per 75 sq. ft. Mix 1 pint to 32 gallons of water per 1200 sq. ft. Water lawn before treating and do not water again for three days.

TICKS: In the yard and in dog kennels spray with 1 ounce per 2 gallons of water or 10 ounces per 20 gallons of water or 1 pint to 32 gallons of water. Spray inside and outside of dog house and other areas where dog sleeps. Put fresh bedding in doghouse after spraying. Spray dogs with same mix as for doghouses to kill ticks on dog. Avoid treatment of puppies under 4 weeks old.

NOTE: Keep children and pets off treated lawn until grass is completely dry.

Liquid Flowable Sevin

A Liquid to be Mixed with Water FOR CONTROL OF MANY INSECTS FEEDING ON FRUIT, VEGETABLES, SHADE TREES, SHRUBS, FLOWERS, AND LAWN GRASSES. KEEP OUT OF REACH OF CHILDREN. CAUTION.

SEE LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT: Carbaryl 90.00%
INERT INGREDIENTS: 10.00%
NET WEIGHT: 1.00 lb (16 oz)
NET VOLUME: 1.00 qt (32 fl oz)

Manufactured by FMC Corporation
P.O. Box 1478
Philadelphia, PA 19102

JAN 12 1989
11:00 AM
1100
1100