

PM-12 869-134 1 1/2

GREEN LIGHT LIQUID FLOWABLE SEVIN<sup>R</sup>

For Control Of Many Insects Feeding On Fruit, Vegetables, Shade Trees, Shrubs, Flowers, And Lawn Grasses

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl methylcarbamate)*.....	28%
INERT INGREDIENTS.....	72%
Total	100%

\*Registered Trademark Union Carbide Corp.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 869-134

EPA Est. No. 869-TX-1

Green Light Co., San Antonio, TX 78217

Net Contents 1 Pt.  
(Back or Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: IF SWALLOWED, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. IF INHALED, remove victim to fresh air and apply respiration if indicated. IF ON SKIN, wash promptly with soap and water. Rinse thoroughly. IF IN EYES, may irritate eyes, nose, throat and skin. Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Subsequent planting of foods and feed crops in treated soil must be made with crops on this or other carbaryl labels.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in a locked storage area out of reach of children and domestic animals. Keep in original container. Do not store near heat or open flame.

PESTICIDE DISPOSAL: Unwanted pesticide material leave in original container. Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

SEE ATTACHED CARD FOR DIRECTIONS FOR USE

Read entire label before using. Use only as directed and follow cautions carefully. SHAKE WELL BEFORE POURING OUT OF BOTTLE.

Dilution: Use 4 teaspoons per 1 gallon of water or 1-1/3 pints per 32 gallons water, except on lawns. Spray all plants thoroughly covering underside and tops of leaves and branches.

SHADE TREES, SHRUBS, FLOWERS: Japanese, blister, flea, and June beetles; boxelder, mealy, lace, and plant bugs; leafhoppers, leafrollers; psyllid, rose aphid and exposed thrips. gypsy moth, bagworms, scale insects; mimosa webworm, tent and puss caterpillars, cicada, orange tortix.

VEGETABLE INSECTS: Tomato, potato, eggplant, and pepper for Colorado potato beetle, flea beetles, lace bugs, tomato hornworm; Beans for Mexican bean beetle, bean leaf beetle, cucumber beetles, Japanese beetle, leafhoppers; Cucumbers, melons, pumpkin, and squash for pickleworm, melonworm, cucumber beetles, squash bugs. Apply 1 gallon spray per 535 sq. ft. Spray may be used within 1 day of harvest.

FRUIT INSECTS: Apples and pears for codling moth, green apple aphid, Japanese beetles, leafroller, bagworm, periodical cicada, San Jose scale crawlers, pear psylla.

NOTE: Sevin<sup>R</sup> Spray will cause thinning of fruit on many apple varieties when applied 10 to 25 days after full bloom. Consult your local County Agricultural Extension Agent for list of varieties affected.

Peaches, plums, apricots, nectarines for codling moth, plum curculio, oriental fruit moth, Japanese beetle, twig borer, periodical cicada, tarnished plant bug, scale crawlers. Spray may be used within 3 days of harvest. Apply spray at petal fall and every 10 to 14 days thereafter until control is achieved.

NOTE: Proper timing of the sprays for control of fruit insects is essential. Consult your local Agricultural Extension Specialists for proper fruit tree spray schedule.

LAWN INSECTS: Chinch bugs, fall armyworms, leafhoppers, sod webworms, ants, earwigs, fleas, mosquitoes. Use 10 ounces per 20 gallons of water per 750 square feet. or 1 ounce per 2 gallons of water per 75 sq. ft. or 1 pint to 32 gallons of water per 1200 sq. ft. Water lawn before treating and do not water again for three days.

TICKS: In the yard and in dog kennels, spray with 1 ounce per 2 gallons of water or 10 ounces per 20 gallons of water or 1 pint to 32 gallons of water. Spray dogs with same mix as for doghouses to kill ticks on dog. Avoid treatment of puppies under 4 weeks old. Do not use on pregnant dogs.

NOTE: Keep children and pets off treated lawn until completely dry. Do not use on Boston Ivy, Virginia Creeper, or Maidenhair Fern.