

174



# GREEN LIGHT

ACCEPTED  
NOV 30 1983  
EPA Reg. No. 869-134

# DIAZINON<sup>®</sup> 5 GRANULES


## FOR CONTROL OF FIRE ANTS

AND CERTAIN OTHER LAWN INSECTS SUCH AS:

CHINCH BUGS, BILLBUGS, WHITE GRUBS such as JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER, as well as BROWN DOG TICKS, LAWN MOTH (SOD WEBWORM), ARMYWORMS, CHIGGERS, LEAFHOPPERS, SOWBUGS, MILLIPEDES, EARWIGS, ANTS, BERMUDA MITES, and FLEAS.



869-139 ©



**ACTIVE INGREDIENT:**  
 O, O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)  
 phosphorothioate\* ..... 5.00%

**INERT INGREDIENTS** ..... 95.00%

**TOTAL** ..... 100.00%

\*Diazinon registered trademark CIBA-GEIGY Corp.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION  
STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

**IF INHALED:** Remove victim to fresh air and apply respiration if indicated.

**IF ON SKIN:** Wash promptly with soap and water. Rinse thoroughly.

**IF IN EYES:** Rinse eyes for at least 15 minutes with water and call a physician if irritation persists.

Note to Physician - If signs of cholinesterase inhibition are present, atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

See other cautions on back of this bag.

Distributed by GREEN LIGHT COMPANY, SAN ANTONIO, TEXAS 78217.

E.P.A. Reg. No. 869-139

E.P.A. Est. No. 869-TX-1

89-137 (3)



**GREEN LIGHT**

**DIAZINON<sup>®</sup> 5  
GRAINULIES**

**FOR CONTROL OF FIRE ANTS**

**AND CERTAIN OTHER LAWN INSECTS SUCH AS:  
CHINCH BUGS, BILLBUGS, WHITE GRUBS such as JAPANESE BEETLE,  
EUROPEAN CHAFER, SOUTHERN CHAFER, as well as BROWN  
DOG TICKS, LAWN MOTH (SOD WEBWORM), ARMYWORMS, CHIGGERS,  
LEAFHOPPERS, SOWBUGS, MILLIPEDES, EARWIGS, ANTS,  
BERMUDA MITES, and FLEAS.**



**PRECAUTIONARY STATEMENTS**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Keep out of reach of children. Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling and before eating or smoking. May be absorbed through skin. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

- STORAGE:** Store in a locked storage area out of reach of children and domestic animals. Store in original container. Do not store near food or feed product.
- PESTICIDE DISPOSAL:** Unwanted pesticide material leave in original container. Securely wrap original container in several layers of newspaper and discard in trash.
- CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

For control of Chinch Bugs, Lawn Moth (sod webworms), Bermuda Mites, Ants, Earwigs, Armyworms, Leafhoppers, Chiggers: On Bermuda, Zoysia, and Bentgrass lawns apply 2 to 3 lbs. Granules per 1,000 sq. ft. On St. Augustine grass lawns apply 3 to 3 1/2 lbs. per 1,000 sq. ft. Water lawn thoroughly and then allow grass to dry before applying Granules. Then water after treatment just enough to get Granules off grass and down to the soil. Do not water lawn for 2 to 3 days after treatment. Repeat application in 7 to 10 days if necessary.

For **WHITE GRUBS** such as Japanese Beetle, European Chafer, Southern Chafer, and Billbugs, apply 2.5 lbs. Granules per 1,000 sq. ft. or 12 lbs. per 4,800 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

**Band Treatment Around Foundation:** Brown Dog Ticks, Crickets, Earwigs, Fleas, Millipedes, and Sowbugs are commonly found around or near foundation of houses as well as in the lawn. Certain of these insects may even enter the house. For control of these insects, apply a five-foot band of Granules next to the foundation of the house. Distribute the Granules uniformly using a rate of 5 oz. for each 5-ft. by 20-ft. area to be treated.

**FIRE ANTS and OTHER ANTS**

**SPOT TREATMENT**

Ants: Apply 1 1/2 teaspoons of Green Light Diazinon 5 Granules around each ant hill.

**Fire Ants:** To aid in the control of fire ants in lawns and other recreational areas, apply 1/2 cup over, and 6 ft. around, each fire ant mound immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

Note: For best results, apply in cool weather, 65-80 degrees F. or in early morning or late evening hour. Treat new mounds as they appear.

**SPREADER SETTING NUMBER PER 1,000 SQ. FT.**

Spreader Make & Model	Regular or Dichondra Lawn	St. Augustine Lawn	White Grubs, Billbugs
	2 lbs	3 lbs	2-1/2 lbs.
Scott	5-1/8	6-1/8	5-5/8
Jackson 18-26 (18")	5	5-3/4	5-1/2
Sears (20")	6-1/4	6-3/4	6-1/2
Salem Micro Set (22")	9-1/4	8-5/8	3-7/8
Cyclone Lawn Spreader	3-1/2	4	3-3/4
Lawn Beauty	6	6-7/8	4-1/2

These settings apply to 20/40 mesh granular carriers. The spreader settings listed above are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting, if necessary, in order to assure correct rate of application.

BERMUDA MITES, and FLEAS

