

419218

JUN - 4 1992

Mr. Jeff Luedke
Green Light Company
P.O. Box 17985
San Antonio, TX 78217

Subject: Amendment - Revise per Diazinon Registration Standard
Green Light Diazinon 5 Granules
EPA Reg. No. 869-139
Your submission received April 23, 1990

Dear Mr. Luedke:

The amendment referred to above, submitted in connection with a registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided the following revisions are made prior to release for shipment under the amended labeling:

- 1) The Diazinon Registration Standard stipulates that diazinon outdoor use products are classified as Restricted Use due to avian and aquatic toxicity unless they are registered for application by homeowners to the home yard and/or home garden or by commercial applicators only to home yards. You can clarify the intended use site for this product by adding to the front panel: "For outdoor use on the home lawn and in the home garden only". A good location for this would be just under the golf course/sod farm prohibition.
- 2) Please note that your newly added claim for control of European Crane Fly would be based on data generated by Chevron Chemical Company. Either submit authorization to use this data or delete the claim.
- 2) The text of the Hazards to Humans and Domestic Animals should be revised as follows to reflect the toxicology data you are relying upon to support reregistration of this product:

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near

food or feed products. Food utensils such as spoons or measuring cups should not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets to occupy treated areas until granules have been watered into the soil and the grass or soil has dried.

- 4) Based on Ciba-Geigy's toxicology data submitted in support of reregistration of their 5% diazinon granular products, and upon which you are relying for reregistration of the subject product, in the Statement of Practical Treatment add "Get medical attention." to the end of the "If on skin" advice.

- 5) Revise the Environmental Hazards to read as follows:

This pesticide is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of granules. For lawns, where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after applying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

- 6) Under the heading "IN GARDENS DIAZINON 5 GRANULES KILLS:", delete the pests "Armyworms", "Carrot rust flies", "Corn borers", and "Rootworms" since the application methods required to control these pests, in-furrow or foliar broadcast, are not being supported for reregistration of granular diazinon products and directions for control of these pests will not appear on the revised label.

- 7) Revise the first sentence of the directions for aiding in control of fire ants to read as follows to indicate the home-use nature of this product: To aid in the control of fire ants in home lawns and other outdoor home recreational areas, apply 1/2 cups over, and 2 ft. around, each fire ant mound.

- 8) For the crop "Beans", delete the directions for control of "Lesser Cornstalk Borers". Ciba-Geigy is not supporting reregistration of granular diazinon applied post-emergence to food crops because of the hazards to birds from exposed granules.
- 9) Revise the variety listing for melons to read as follows:

MELONS (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, and Hybrids of these) and WATERMELONS
- 10) For the crop "Onions", delete "by disking, harrowing, or other suitable means". This instruction was a carryover from an agricultural product label and is not appropriate to a homeowner-use product.
- 11) For the crop "Potatoes", delete the "Note" about hand-harvested commercially-grown potatoes, since this is not appropriate to a product whose crop uses are limited to application to the home garden.
- 12) For the crop "Sweet Corn" delete the directions for use for control of "European Corn Borers", "Fall Armyworms", and "Southwestern Corn Borers". Also, delete the "NOTES: Corn may be picked...application". Ciba-Geigy is not supporting reregistration of granular diazinon applied post-emergence to food crops because of the hazards to birds from exposed granules.
- 13) For the crop "Sweet Potatoes", delete the special directions for post-emergence use in the Southeastern U. S.. Ciba-Geigy is not supporting reregistration of granular diazinon applied post-emergence to food crops because of the hazards to birds from exposed granules.

A stamped copy of the label is enclosed for your records. Please submit five copies of the revised amended label prior to release for shipment under the amended label. Labeling released for shipment after the August 31, 1992 deadline imposed by the Diazinon Registration Standard should be the revised amended label.

Sincerely yours,

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (H7505C)

GREEN LIGHT DIAZINON 5 GRANULES

Lawn & Garden Insect Control

Treats up to
4,000 sq. ft.

Kills Ants, Armyworms, Bermudagrass Mites, Billbugs, Brown Dog Ticks, Chiggers, Chinch Bugs, Crickets, Cutworms, Earwigs, European Cranefly (larvae), Fire Ants (aids in control), Fleas, Millipedes, Root Maggots, Sod Webworms (Lawn Moths), Sowbugs, White Grubs of: Japanese Beetle; European Chafer; Southern Chafer, Wireworms, and others

This Product Must Not Be Used On Golf Courses And Sod Farms

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)	
phosphorothioate.....	5%
INERT INGREDIENTS.....	95%
	Total 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Additional Precautionary Statements On Back Of Bag

EPA Reg. No. 869-139

EPA Est. No. 869-TX-1

Net Wt. 8 lbs.

Green Light Company, San Antonio, TX 78217

ACCEPTED
with COMMENTS
In EPA Letter Data
JUN - 4 1992

Under the Federal Insecticide
Fungicide and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
869-139

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before eating or smoking. May cause contact sensitization following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: call a physician immediately. Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. IF INHALED: remove victim to fresh air and apply artificial respiration, if indicated. IF ON SKIN: wash promptly with soap and water. Rinse thoroughly. IF IN EYES: rinse eyes with plenty of water and call a physician immediately.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS: This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of $\frac{1}{4}$ inch of water immediately after applying this product. Stop irrigation before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries.

Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Runoff may be hazardous to aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning or equipment or disposal of equipment wash waters.

In Lawns Diazinon 5 Granules Kills:

Ants	Crickets	Sod webworms
Armyworms	Cutworms	Sowbugs
Bermudagrass mites	Digger wasps	Springtails
Billbugs	Earwigs	White grubs of:
Brown dog ticks	Fire ants*	• Japanese beetle
Chiggers	Fleas	• European chafer
Chinch bugs	Lawn moths	• Southern chafer
Clover mites	Leafhoppers	Mole Crickets
European Crane fly (larvae)	Millipedes	* Aids in control

In Gardens Diazinon 5 Granules Kills:

Armyworms	Mole crickets	Rootworms
Carrot rust flies	Onion maggots	Vinegar flies
Corn borers	Root maggots	Wireworms
Cutworms		

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product.

Conditions of Sale and Warranty

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of G.L. Co. or the Seller. All such risks shall be assumed by the Buyer.

Green Light warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. G.L. Co. makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Green Light or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. G.L. Co. and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Green Light.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Overall Lawn Application

Diazinon 5 G controls the following lawn insects on both grass and dicotyledon lawns:

Ants • Armyworms • Bermudagrass mites • Brown dog ticks • Chiggers • Chinch bugs • Clover mites • Crickets • Cutworms • Earwigs • Fleas • Lawn moths (sod webworms) • Leafhoppers • Millipedes • Sowbugs • Springtails (Collembola) • European Crane fly (larvae), Mole Crickets.

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds. Avoid overlapping treated areas. If waterfowl, i.e., ducks, geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 1/4 inch water immediately after application, however, stop watering before puddling occurs.

Apply granules with a fertilizer spreader. See spreader setting chart below. Apply when grass is dry. Repeat as necessary.

For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10').

Billbugs; White grubs of Japanese beetle, European chafer, Southern chafer, *Ataenius spretulus* (dung beetle): Apply 2 lbs. per 1,000 sq. ft. of lawn. Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or early Summer when damage is first noticed. Water grass thoroughly with 1/4-1/2 in. of water after application. Treat again in late Summer or early Fall if a second generation occurs.

Hyperodes weevils

Apply 2 lbs. per 1,000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

LAWN SPREADER SETTING NO. (per 1000 sq. ft.)

Spreader Make and Model	Lawns/2 lbs.
Central Imperial Mark No. 1662-G	6 1/2
Central Suburbia Lawn Spreader No. 1622-0	6 1/2
Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader Model 3000	4 1/2
ORTHO Broadcast Spreader Model 4000	4 1/2
J. C. Penney Model 6005	7 1/4
J. C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF-4	4
Sears Craftsman Model 671.19198	5

Spot Treatment Lawns

Ants: Apply 1½ teaspoons granules over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other recreational areas, apply ½ cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary.

Notes: For best results, apply in cool weather, 65-80°F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

Digger wasps: Apply 1½ teaspoons granules in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses.

- | | | |
|-----------------|----------|--------------------------|
| Ants | Crickets | Millipedes |
| Brown dog ticks | Earwigs | Sowbugs (Pillbugs) |
| Clover mites | Fleas | Springtails (Collembola) |

Apply a 5 foot band next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. Repeat as necessary.

Vegetables - Insect Control

Diazinon 5 G controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Unless directions under "Timing/Special Directions" state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

Insect	Depth of Mixing or Incorporation into Soil
Surface cutworms *	2-3 inches
Subterranean cutworms *	3-6 inches
Mole crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

* If type of cutworm cannot be identified, mix into soil 4-6 inches deep.

CROP	PEST	RATE	TIMING/SPECIAL DIRECTIONS
Beans (Pole, Snap or Lima) (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Lesser Cornstalk Borers	3½-7 oz. per 500 sq. ft.	Apply in a 6-inch band centered over the row when plants are ½-1 inch tall.
NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc.			
Beets, Red (Table)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Broccoli	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Brussels Sprouts	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cabbage	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.

Carrots	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cauliflower	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Celery	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Collards	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cucumber	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Endive (Escarote)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Kale	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.

Lettuce	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, & Hybrids of these and watermelons)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Mustard	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Onions (green and bulb)	Onion Maggots	7-14 oz. per 500 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil by disking, harrowing, or other suitable means. NOTE: Diazinon may not control organo-phosphate-resistant onion maggots.
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Parsley	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Peas (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
NOTE: Do not use on dried pea varieties such as dried blackeyed peas.			
Peppers	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Potatoes	Cutworms (Surface & Subterranean)	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
NOTE: Do not use on commercially-grown potatoes which will be hand-harvested.			

Radishes	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Spinach	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Squash, Summer and Winter	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Sweet Corn	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Seed Corn Maggots	7-14 oz. per 500 sq. ft.	Broadcast just before planting. Mix into soil 2 inches.
	European Corn Borers	3½-7 oz. per 500 sq. ft.	When 75% of the plants show first-generation borers feeding, apply over the corn so granules fall into whorls. Make a second application 7 days later if needed. For second-generation borers, apply as above when there is an average of 1 egg mass per plant. Do not make more than 3 applications per season.
	Fall Armyworms	3½-7 oz. per 500 sq. ft.	Apply over corn so granules fall into whorls when larvae first appear. If necessary, application may be repeated twice at 7-day intervals. Do not make more than 3 applications per season.
	Southwestern Corn Borers	3½-7 oz. per 500 sq. ft.	7-day intervals. Do not make more than 3 applications per season.
NOTES: Corn may be picked immediately after last application. Do not feed fodder to livestock for 10 days following last application.			

Sweet Potatoes	Wireworms Flea Beetles	10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting Southeastern U.S. - Broadcast 10½ oz. per 500 sq. ft. over the tops of vines when roots begin to enlarge, but do not mix into soil.
Swiss Chard	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Tomatoes	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Turnips	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.

Storage and Disposal

Storage: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse empty bag. Discard bag in trash. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.