

CAUTION: KEEP OUT OF REACH OF CHILDREN. HARMFUL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN. MAY BE IRRITATING TO EYES AND SKIN. COMBUSTIBLE.

Avoid skin and eye contact. Do not smoke while using. Wash thoroughly after handling. Flush contaminated eyes with plenty of water and get medical attention. Avoid breathing mist of vapor. Wash contaminated clothing before re-use.

NOTE TO PHYSICIAN:

Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL CAUTIONS:

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Agricultural Extension Service. Apply this product only as specified on this label. Keep this product in original container.

Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not spray vegetable gardens or other plants used for food purposes. Keep away from food, feed stuffs, and domestic water supplies. Keep container closed.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. Handle concentrate in a ventilated area.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

DIRECTIONS FOR USE

For control of chinchbugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs, and grasshoppers infesting lawns and turfs. Wet grass down thoroughly before spraying with this emulsifiable concentrate. Dilute spray at the rate of 2 tablespoonful per gallon of water to treat 50 square feet or 1 gallon per 125 to 150 gallons of water and apply over 6,600 square foot area. When using a lawn hose-on-sprayer (15-gallon to 20-gallon sprayer), fill quart jar to 20-ounce mark with spray and finish filling jar to top with water. Then apply this mixture over 1,000 square feet of lawn grass. Ten steps in one direction and then in another direction is approximately 1,000 square feet. Repeat application if chinch bug damage reoccurs or when pests reappear.


NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building and also the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Dilute spray at the same rate as above. Apply as a coarse spray to thoroughly and uniformly wet the band area.

To control hyperodes (grass) weevils, use 2 gallons spray per 125 to 150 gallons water and apply over 6,600 square foot area, or 4 tablespoonfuls per gallon of water to treat 50 square feet. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

CAUTION: Keep children and pets off treated areas until spray has dried.

ORNAMENTALS: To control aphids, bagworms, grasshoppers, mites, spittlebugs, whiteflies: Dilute 2 ounces (4 tablespoons) in 1 gallon of water. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10-day intervals if needed.

NOTE: Do not use on azaleas, camellias, hydrangeas, petas, roses, or variegated ivy because of possible injury to the plants and ants.



869-149
12-10-75

...HARMFUL
...SKIN. MAY
...E.

...using. Wash
...eyes with
...breathing
...before re-use.

Treat sympto-

...wildlife. Keep
...runoff is
...conditions favor

...direct treat-
...ation may be
...Service. Apply
...Keep this prod-

...equipment or dis-

...plants used for
...ed stuffs, and
...sed.

...HEAT OR OPEN
...rea.

...y perforating or
...rom water

DIRECTIONS FOR USE

For control of chinchbugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs, and grasshoppers infesting lawns and turfs. Wet grass down thoroughly before spraying with this emulsifiable concentrate. Dilute spray at the rate of 2 tablespoonful per gallon of water to treat 50 square feet or 1 gallon per 125 to 150 gallons of water and apply over 6,600 square foot area. When using a lawn hose-on-sprayer (15-gallon to 20-gallon sprayer), fill quart jar to 20-ounce mark with spray and finish filling jar to top with water. Then apply this mixture over 1,000 square feet of lawn grass. Ten steps in one direction and then in another direction is approximately 1,000 square feet. Repeat application if chinch bug damage reoccurs or when pests reappear.


NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building and also the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Dilute spray at the same rate as above. Apply as a coarse spray to thoroughly and uniformly wet the band area.

To control hyperodes (stargrass) weevils, use 2 gallons spray per 125 to 150 gallons water and apply over 6,600 square foot area, or 4 tablespoonfuls per gallon of water to treat 50 square feet. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

CAUTION: Keep children and pets off treated areas until spray has dried.

ORNAMENTALS: To control aphids, bagworms, grasshoppers, mites, spittlebugs, whiteflies: Dilute 2 ounces (4 tablespoons) in 1 gallon of water. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10-day intervals if needed.

NOTE: Do not use on azaleas, camellias, petas, roses, or variegated ivy because of possible injury to the plants.



ACTIVE INGREDIENTS:

Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate]* 2.0%
Xylene 88.0%

INERT INGREDIENTS 10.0%
Total 100.0%

Contains 2.5 ounces Chlorpyrifos per gallon

*U. S. Patent No. 3,244,586

*Dursban reg. trademark The Dow Chem. Co.

READ INSTRUCTIONS CAREFULLY
KEEP OUT OF REACH OF CHILDREN

Read complete precautions on back panel.

GREEN TICK COMPANY San Antonio, Texas 78215

E.P.A. Est. No. 869-TX-1 E.P.A. Reg. No. 869-149-AA

NET CONTENTS ONE PINT

869-149
12-10-75

...EN. HARMFUL
...ON SKIN. MAY
...IBLE.

...e using. Wash
...ted eyes with
...void breathing
...before re-use.

Treat sympto-

...wildlife. Keep
...ere runoff is
...onditions favor

...to direct treat-
...ation may be
...Service. Apply
...Keep this prod-

...ipment, or dis-

...plants used for
...ed stuffs, and
...sed.

...HEAT OR OPEN
...rea.

...y perforating or
...rom water

DIRECTIONS FOR USE

For control of chinchbugs, sod webworms (lawn moths), ants, brown dog ticks, crickets, cutworms, earwigs, and grasshoppers infesting lawns and turfs. Wet grass down thoroughly before spraying with this emulsifiable concentrate. Dilute spray at the rate of 2 tablespoonsful per gallon of water to treat 50 square feet or 1 gallon per 125 to 150 gallons of water and apply over 6,600 square foot area. When using a lawn hose-on-sprayer (15-gallon to 20-gallon sprayer), fill quart jar to 20-ounce mark with spray and finish filling jar to top with water. Then apply this mixture over 1,000 square feet of lawn grass. Ten steps in one direction and eleven in another direction is approximately 1,000 square feet. Repeat application if chinch bug damage reoccurs or when pests reappear.

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building and also the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Dilute spray at the same rate as above. Apply as a coarse spray to thoroughly and uniformly wet the band area.

To control hyperodes (turfgrass) weevils, use 2 gallons spray per 125 to 150 gallons water and apply over 6,600 square foot area, or 4 tablespoonfuls per gallon of water to treat 50 square feet. Spray suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

CAUTION: Keep children and pets off treated areas until spray has dried.

ORNAMENTALS: To control aphids, bagworms, grasshoppers, mites, spittlebugs, whiteflies: Dilute 2 ounces (4 tablespoons) in 1 gallon of water. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10-day intervals if needed.

NOTE: Do not use on azaleas, camellias, petas, roses, or variegated ivy because of possible injury to the ants.

867-149
12-10-75



Controls Pests on Turf: Chinch Bugs, sod webworms, ants, brown dog ticks, crickets, cutworms, earwigs, grasshoppers, hyperodes weevils. Controls pests on Ornamental Plants: Aphids, bagworms, grasshoppers, mealybugs, mites, spittlebugs, whiteflies.

ACTIVE INGREDIENTS:

Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate]* 2.0%
Xylene 88.0%

INERT INGREDIENTS 10.0%
Total 100.0%

Contains 2.5 ounces Chlorpyrifos per gallon

*U. S. Patent No. 3,244,586

*Dursban reg. trademark The Dow Chem. Co.

**READ ENTIRE LABEL BEFORE USING.
KEEP OUT OF REACH OF CHILDREN**

CAUTION

Read complete precautions on back panel.

GREEN LIGHT COMPANY San Antonio, Texas 78246

E.P.A. Est. No. 869-TX-1 E.P.A. Reg. No. 869-149-AA

NET CONTENTS ONE PINT