

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)**

CAUTION

Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after using. Harmful if swallowed. Avoid contamination of feed and foodstuffs. Keep out of reach of children and domestic animals.
PHYSICIAN: Atropine Sulfate is antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. This product is highly toxic to bees exposed to direct treatment or residue on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Leave in original container until used. Do not store near heat or open flame. **STORE OUT OF REACH OF CHILDREN.**

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Proper timing, application rates, and repeated sprayings are necessary for fruit insect control. Always consult your local Agricultural Extension Specialist for fruit tree spray schedule.

Mix one pint Fruit and Vegetable Spray in 48 gallons of water or one ounce (2 tablespoons) in 3 gallons of water to spray fruit trees.

NOTE: Do not spray apples, pears, grapes, or cherries within 3 days of harvest.

APPLES AND PEARS: Codling moth, plum curculio, spray when 3/4 petal blossoms have fallen and repeat at 10-day intervals. Usually 4 or 5 sprayings are necessary to protect fruit from larvae. Spray when woolly and green aphids, red-banded leaf rollers and tent caterpillars first appear on twigs and leaves. Repeat at 10-day intervals until pests are controlled.

GRAPES: Spray when leaf hoppers or mealy bugs appear. Repeat at 10-day intervals as necessary. Do not spray more than 250 gallons per acre.

CHERRIES: Spray when black cherry aphids, Japanese beetles, or cherry fruit fly appear and repeat at 10-day intervals as necessary. Do not spray more than 700 gallons per acre.

NOTE: Do not spray peaches, quince, plums, or apricots within 7 days of fruit harvest. Spray may injure foliage on some varieties of apples, grapes, cherries.

PEACHES, QUINCE, PLUMS: For control of plum curculio, oriental fruit moth, spray when half the petal blossoms have fallen and repeat at 10-day intervals. Usually 4 or 5 sprayings are necessary to protect fruit from larvae. Spray when aphids, Japanese beetles first appear. Repeat at 10-day intervals if necessary.

APRICOTS: For codling moth, spray when half the petal blossoms have fallen and repeat 4 or 5 times at 10-day intervals.

CITRUS: Red scale, black scale, purple scale, soft scale, Mediterranean fruit fly, aphids, thrips, spray with 1 to 1 1/2 quarts per 100 gallons of water or 2 or 3 tablespoons per 6 gallons of water. Spray when insects first appear and repeat applications at 10-day intervals until pests are controlled. Make no spray application within 7 days of harvest.

STRAWBERRIES: Spray when potato leafhoppers, strawberry root weevil first appear and repeat at 10-day intervals as needed. Do not apply spray within 3 days of harvest.

VEGETABLE INSECTS: Mix 1 pint Fruit and Vegetable Spray in 48 gallons of water or 1 ounce (2 tablespoons) in 3 gallons of water. Spray when insects first appear and repeat at 10-day intervals as needed to control pests.

TOMATOES: for aphids, spider mites, BEANS, SQUASH, CUCUMBERS for aphids, leafhoppers, cucumber beetles. **NOTE:** Do not spray these vegetables within 1 day of harvest. **EGGPLANT, PEPPERS:** for aphids, lace bugs.

NOTE: Do not spray within 3 days of harvest.


MUSTARD GREENS, TURNIPS, SPINACH: for aphids, leafhoppers, flea beetles.

NOTE: Do not spray within 7 days of harvest. Do not use this spray on any plants or in any way other than as directed on this label.



**GREEN LIGHT
TOMATO & VEGETABLE
SPRAY CONCENTRATE**

For Home Fruit Trees, Home Vegetable Gardens
Controls Fruit and Vegetable Insects Listed Below
READ ENTIRE LABEL BEFORE USING.



ACTIVE INGREDIENTS	
*Methionin	39.50%
Xylene	39.00%
INERT INGREDIENTS	
	11.00%
Total	100.00%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate. Derived from 95% technical Methionin.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF INHALED: Remove victim to fresh air and apply respiration if indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.

IF IN EYES: Rinse eyes for at least 15 minutes with water and call a physician immediately.

SEE ADDITIONAL CAUTIONS ON BACK OF PANEL.


E.P.A. Est. No. 888-TX-1

E.P.A. Reg. No. 888-152

Distributed by GREEN LIGHT COMPANY, San Antonio, Texas 78217

NET CONTENTS 1/2 Gallon

1/2 gal



P.M. 16 869-152